



PCT

WORLD INTELLECTUAL PROPERTY
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER T.

WO 9605303A1

(51) International Patent Classification 6 : C12N 15/12, C07K 14/705, A61K 39/395, C12N 15/11, A61K 38/17, C07K 16/28, G01N 33/68	A1	(11) International Publication Number: WO 96/05303 (43) International Publication Date: 22 February 1996 (22.02.96)
(21) International Application Number: PCT/US95/09636 (22) International Filing Date: 9 August 1995 (09.08.95) (30) Priority Data: 289,699 12 August 1994 (12.08.94) US (71) Applicant: OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 N.E. Thirteenth Street, Oklahoma City, OK 73104 (US). (72) Inventors: FUKUDOME, Kenji; 13125 West Park Place, Oklahoma City, OK 73142 (US). ESMON, Charles, T.; 5800 North Stonewall, Oklahoma City, OK 73111 (US). (74) Agent: PABST, Patrea, L.; Arnall Golden & Gregory, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: CLONING AND REGULATION OF AN ENDOTHELIAL CELL PROTEIN C/ACTIVATED PROTEIN C RECEPTOR (57) Abstract <p>Human protein C and activated protein C were shown to bind to endothelium specifically, selectively and saturably ($K_d=30$ nM, 7000 sites per cell) in a Ca^{2+} dependent fashion. Expression cloning revealed a 1.3 kb CDNA that coded for a novel type I transmembrane glycoprotein capable of binding protein C. This protein appears to be a member of the CD1/MHC superfamily. Like thrombomodulin, the receptor involved in protein C activation, the endothelial cell protein C receptor (EPCR) function and message are both down regulated by exposure of endothelium to TNF. Identification of EPCR as a member of the CD1/MHC superfamily provides insights into the role of protein C in regulating the inflammatory response, and determination of methods for pharmaceutical use in manipulating the inflammatory response.</p>		

[illegible]

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

BNSDOCID: <WO 9605303A1 I_>

CLONING AND REGULATION OF AN ENDOTHELIAL CELL
PROTEIN C/ACTIVATED PROTEIN C RECEPTOR

Background of the Invention

The present invention is generally in the area of cloning, expression, and regulation of an endothelial cell protein C/activated protein C
5 receptor.

Protein C plays a major role in the regulation of blood coagulation. Patients deficient in protein C usually exhibit life threatening thrombotic-complications in infancy
10 (Seligsohn et al., (1984) N. Engl. J. Med. 310, 559-562; Esmon, (1992) Trends Cardiovasc. Med. 2, 214-220) that are corrected by protein C administration (Dreyfus et al., (1991) N. Engl. J. Med. 325, 1565-1568). In addition, activated
15 protein C (APC) can prevent the lethal effects of *E. coli* in baboon models of gram negative sepsis (Taylor et al., (1987) J. Clin. Invest. 79; U.S. Patent No. 5,009,889 to Taylor and Esmon) and preliminary clinical results suggest that protein C
20 is effective in treating certain forms of human septic shock (Gerson et al., (1993) Pediatrics 91, 418-422). These results suggest that protein C may both control coagulation and influence inflammation. Indeed, inhibition of protein S, an
25 important component of the protein C pathway, exacerbates the response of primates to sublethal levels of *E. coli* and augments the appearance of TNF in the circulation (Taylor et al., (1991) Blood 78, 357-363). The mechanisms involved in
30 controlling the inflammatory response remain unknown.

Protein C is activated when thrombin, the terminal enzyme of the coagulation system, binds to an endothelial cell surface protein, thrombomodulin
35 (Esmon, (1989) J. Biol. Chem. 264, 4743-4746; Dittman and Majerus, (1990) Blood 75, 329-336;

Dittman, (1991) Trends Cardiovasc. Med. 1, 331-336). In cell culture, thrombomodulin transcription is blocked by exposure of endothelial cells to tumor necrosis factor (TNF) (Conway and Rosenberg, (1988) Mol. Cell. Biol. 8, 5588-5592) and thrombomodulin activity and antigen are subsequently internalized and degraded (Lentz et al., (1991) Blood 77, 543-550, Moore, K.L., et.al., (1989) Blood 73, 159-165). In addition, C4bBP, a regulatory protein of the complement system, binds protein S to form a complex that is functionally inactive in supporting APC anticoagulant activity in vitro (Dahlbäck, (1986) J. Biol. Chem. 261, 12022-12027) and in vivo (Taylor, et al., 1991). Furthermore, C4bBP behaves as an acute phase reactant (Dahlbäck, (1991) Thromb. Haemostas. 66, 49-61). Thus, proteins of this pathway not only appear to regulate inflammation, but they also interact with components that regulate inflammation, and they themselves are subject to down regulation by inflammatory mediators.

Given the central role of the protein C pathway in regulating the host response to inflammation and the critical role of the pathway in controlling blood coagulation, it is important to identify and characterize all of the components that interact with the system. This is especially true since the molecular basis of the anti-inflammatory effects of the protein C pathway components have yet to be elucidated at the molecular level.

It is therefore an object of the present invention to provide a cellular receptor for protein C and activated protein C.

It is a further object of the present invention to provide nucleotide sequences encoding

the cellular receptor and amino acid characterization of the Receptor which allows expression of recombinant native and modified forms of the receptor.

5 It is another object of the present invention to provide methods of modulating the inflammatory response involving protein C and activated protein C.

Summary of the Invention

10 An endothelial cell protein C binding protein (referred to herein as "EPCR") has been cloned and characterized. The protein is predicted to consist of 238 amino acids, which includes a 15 amino acid signal sequence at the N-terminus, and a 23 amino acid transmembrane region which characterizes the receptor as a type 1

transmembrane protein. The protein binds with high affinity to both protein C and activated protein C ($K_d=30$ nM) and is calcium dependent. The message and binding function of the receptor are both down regulated by cytokines such as TNF.

20 These results identify a new member of a complex pathway that, like other members of the pathway, is subject to regulation by inflammatory cytokines, and can therefore be used to modulate inflammatory reactions in which protein C or activated protein C is involved. Inhibition of the inflammatory response can be obtained by infusing soluble EPCR. Alternatively, localizing EPCR to surfaces in contact with blood will render the surfaces anticoagulant by virtue of the ability of EPCR to bind and concentrate the anticoagulant activated protein C at the surface. Alternatively, the function of EPCR can be enhanced by overexpressing the EPCR in endothelium that could

35

be used to coat vascular grafts in patients with vascular disease or on stents in cardiac patients.

Brief Description of the Drawings

Figures 1A, 1B and 1C are flow cytometric analyses of F1-APC (fluorescent labelled activated protein C) binding to HUVEC (human umbilical vein endothelial cells). Figure 1A is a graph of cell number versus log of fluorescence intensity, demonstrating F1-APC binding to HUVEC. HUVEC (1×10^5) were incubated at room temperature without (dotted line) or with 160 nM of F1-APC (solid line) in the presence of 1.3 mM CaCl_2 . After washing, bound APC was analyzed by flow cytometry. Figure 1B is a graph of fluorescence intensity versus F1-APC concentration (nM) demonstrating the concentration dependence of F1-APC binding to HUVEC. HUVEC were incubated with F1-APC in the absence (open circles) or presence of 1.3 mM CaCl_2 (closed circles) and binding was measured as in A. Mean channel fluorescence intensity is plotted for each F1-APC concentration (between 0 and 800 nM). Figure 1C is a graph of the percent of mean fluorescence versus unlabeled protein concentration ($\mu\text{g/ml}$), demonstrating the effects of unlabeled proteins on F1-APC binding to HUVEC. F1-APC binding to HUVEC was carried out in the presence of the indicated concentrations (between 0 and 100 $\mu\text{g/ml}$) of unlabeled APC, protein C, protein S, factor X and Xa or recombinant Gla-domainless protein C (rGDPC).

Figures 2A, 2B, 2C and 2D are graphs of ^{125}I -APC Binding to HUVEC Monolayers. Figure 2A is a graph of the bound APC ($\text{cpm} \times 10^{-3}$) versus time (min), showing the time course of ^{125}I -APC binding to HUVEC. HUVEC monolayers (1.2×10^5 cells) were incubated at 4°C with 32 nM (filled squares) or 8

nM (open squares). ^{125}I -APC. At the indicated times, cells were washed and bound radioactivity was measured. Figure 2B is a graph of bound APC (cpm $\times 10^{-3}$) versus unlabeled protein (nM) demonstrating the effects of unlabeled APC and rGDPC on ^{125}I -APC binding to HUVEC. HUVEC were incubated at 4°C for one hour with ^{125}I -labeled APC in the presence of the indicated concentrations (between 01 and approximately 1000 nM) of unlabeled APC (open circles) or rGDPC (closed circles). After washing, bound radioactivity was measured. Figure 2C is a graph of bound APC (fmol/well) versus free APC (nM) demonstrating the concentration dependence of ^{125}I -APC binding to HUVEC. Monolayers of HUVEC were incubated with the concentrations of ^{125}I -APC indicated as described above. Specific binding was determined as described below. Figure 2D is a Scatchard analysis of ^{125}I -APC binding to HUVEC. Each value was calculated from the data shown in Figure 2C.

Figures 3A and 3B are flow cytometric analyses of F1-APC binding to 293T cells transfected with a cDNA clone of EPCR. Cells were transfected with a clone EPCR/pEF-BOS or pEF-BOS (negative control) by the calcium/phosphate method. After 24 h, cells were harvested and F1-APC binding was performed in the absence (dotted lines) or presence of 1.3 mM CaCl_2 (solid lines).

Figures 4a-4c are the predicted protein structure of EPCR based on nucleotide sequence (SEQ ID NO. 1), predicted amino acid sequence (SEQ ID NO. 2) and a hydropathy plot of EPCR. The signal sequence and transmembrane region are indicated with the solid bars.

Figures 5a-5b are a comparison of the amino acid sequence of EPCR to the amino acid sequences of other members of the CD1 family and CCD41. The EPCR sequence is shown in the first line and

compared to murine CCD41 (second line), human CD1d (third line) and murine CD1.2 (fourth line).

Identities with EPCR are indicated by open boxes.

Residues that are conserved between EPCR and all of the human CD1 family members are indicated by a double asterisk. Residues shared with one or more members of the CD1 family are indicated by a single asterisk.

Figure 6 is a comparison of the amino acid sequence of human EPCR (first line) to the amino acid sequence of murine EPCR (second line). Identities are indicated by lines. Similarities are indicated with dots.

Detailed Description of the Invention

I. Cloning and Characterization of EPCR.

Human protein C and activated protein C are shown to bind to endothelium specifically, selectively and saturably ($K_d = 30$ nM, 7000 sites per cell) in a Ca^{2+} dependent fashion. FL-APC binding to various human cell lines were examined, and found that the binding was HUVEC specific. A human kidney cell line transformed with SV40 large T antigen, 293T cells, expressed very few of these binding sites. A HUVEC cDNA library was constructed using the powerful mammalian expression vector, pEF-BOS (Mizushima and Nagata, (1990) Nucleic Acids Res. 18, 5322). Plasmid DNA was prepared from subpools of independent colonies (2,500 colonies per pool), and transfected into 293T cells, using the method of Kaisho et al., (1994) Proc. Natl. Acad. Sci. (USA) 91, 5325. FL-APC binding was analyzed on a flow cytometer. One of eight subpools gave a positive signal. This subpool was divided into 20 subpools and rescreened. After three rounds of enrichment, one positive clone, EPCR-1, was isolated. EPCR-1

SUBSTITUTE SHEET (RULE 26)

carries a 1.3 kb insert. When transfected into 293T cells, this clone was capable of expressing the calcium-dependent binding site for FL-APC on the 293T cell surface.

5 Expression cloning revealed a 1.3 kb cDNA that coded for a type I transmembrane glycoprotein capable of binding protein C. This protein appears to be a member of the CD1/MHC superfamily. Like thrombomodulin, the receptor involved in protein C
10 activation, the endothelial cell protein C receptor (EPCR) function and message are both down regulated by exposure of endothelium to TNF. Identification of EPCR as a member of the CD1/MHC superfamily provides insights into the role of this receptor
15 for protein C in regulating the inflammatory response.

Materials and Methods

Protein Preparation

Human protein C (Esmon et al., (1993) Meths. Enzymol. 222, 359-385), APC (Esmon et al.,
20 1993), recombinant gla domainless protein C (rGDPC) (Rezaie et al., (1992) J. Biol. Chem. 267, 11701-11704), protein S (Taylor et al., 1991), factor X and factor Xa (Le Bonniec et al., (1992) J. Biol. Chem. 267, 6970-6976) were prepared as
25 described in the cited publications.

Selective labeling of the active site of APC with fluorescein was performed by the method of Bock (Böck, P.E. (1988) Biochemistry 27,
30 6633-6639). In brief, N^ε-[[(acetylthio)acetyl]-D-Phe-Pro-Arg-CH₂Cl (200 μM) was reacted with 40 μM APC for 1 hour at room temperature. After dialysis, the covalently modified APC was incubated at room temperature for
35 one hour with 200 μM 5-(iodoacetamido)fluorescein (Molecular Probes). Free fluorescein was removed by gel filtration on a PD-10 column (Pharmacia).

With this method, each molecule of fluoresceinated APC (FI-APC) contains a single dye at the active site and hence all of the fluorescent molecules behave identically.

- 5 Iodogen (Pierce) was used to radiolabel APC with Na¹²⁵I (Amersham) according to the manufacture's protocol in the presence of 5 mM CaCl₂. Free ¹²⁵I was removed by gel filtration on a PD-10 column. The specific activity of the ¹²⁵I-APC
10 was 1×10^4 cpm/ng protein.

Cell Culture

- Human umbilical vein endothelial cells (HUVEC) were isolated from fresh umbilical-cords by collagenase treatment and cultured in medium 199
15 containing 15% fetal bovine serum, 10 µg/ml heparin, and 0.5% endothelial cell growth supplement prepared from bovine brain extract (Maciag et al., (1979) Proc. Natl. Acad. Sci. (USA) 76, 5674-5678). HOS (ATCC CRL 1543), HEP-2 (ATCC
20 CCL 23) and 293 cells (ATCC CRL 1573) transformed with SV40 large T antigen (293T, a gift from Dr. Kenji Oritani) were maintained in Earl's MEM supplemented with 10% fetal bovine serum. The human lymphocyte cell lines, Jurkat, MOLT3 (ATCC
25 CRL 1552), Jijoye (ATCC CCL 87), Raji (ATCC CCL 86), U-937 (ATCC CRL 1593), HL-60 (ATCC CCL 240), and HEL (ATCC TIB 180), were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum.

30 Flow Cytometric Analysis of FI-APC Binding to Cells

- Adherent cells were harvested by incubation at 37°C for 5 min in phosphate buffered saline (PBS) containing 0.02% EDTA. Cells were
35 washed twice with EDTA/PBS and then once with Hank's balanced salt solution (HBSS). They were resuspended in HBSS containing 1% bovine serum albumin (BSA) and 0.02% sodium azide (binding

buffer). Cells (1×10^5) were incubated at room temperature for 45 min with F1-APC in the dark. After washing, they were resuspended in the binding buffer containing 0.5 $\mu\text{g/ml}$ of propidium iodide. Bound F1-APC was analyzed on a flow cytometer, FACScan (Becton Dickinson). Living cells were gated on a dot plot display of forward-scatter (FSC) versus fluorescence-2 (FL2), and F1-APC binding was detected on the fluorescence-1 (FL-1) channel. All experiments were performed in duplicate.

¹²⁵I-APC binding to HUVEC

Monolayers of HUVEC in 24-well microplates (Costar) (1×10^5 cells per well) were washed twice with EDTA/PBS and once with ice-cold HBSS. Cells were then incubated at 4°C for one hour in the binding buffer with ¹²⁵I-APC. After washing three times with ice-cold HBSS, cells were released with the EDTA buffer, and the bound radioactivity was measured in a gamma counter (Isodata 500). To determine non-specific, calcium-independent adsorption of radioactivity, the cells were washed with EDTA/PBS and residual radioactivity in the cell pellet was measured. Non-specific binding of radioactivity was consistently less than 5% of the specific binding. The data was analyzed using the Enzfitter program (Elsevier Biosoft, Cambridge, U.K.).

Construction of HUVEC CDNA Library

Poly-A RNA was isolated from HUVEC (1×10^6 cells) using the FastTrack™ mRNA isolation kit (Invitrogen). cDNA was synthesized from 3 μg of poly-A RNA using a Librarian™ I kit (Invitrogen). A BstX I adaptor was ligated, double stranded cDNA was fractionated by agarose gel electrophoresis, and cDNA longer than 700 bp was ligated into a mammalian expression vector, pEF-BOS (Mizushima and

Nagata, 1990; this vector was a kind gift from Dr. S. Nagata). The construct was transfected into *E. coli* DH10B by electroporation (Bio-Rad Gene Pulser™). The library consisted of 8×10^6 independent colonies with an average size of 2.0 kb.

Expression Cloning and Sequence Analysis

Approximately 2×10^4 independent colonies were divided into eight subpools (each containing 2,500 independent colonies) and plasmid DNA was prepared from each subpool. Sub-confluent 293T cells in 24-well microplates were transfected with 1 μ g of the DNA by the calcium/phosphate method (Graham and Van Der Eb, (1973) *Virology* 52, 456-467). After 20 hours, the medium was changed, and culture was continued for another 24 hours. The subpools were screened for F1-APC binding by FACS analysis as described above. The positive library pool was then divided into 20 new pools and rescreened. After three rounds of screening, 96 individual clones were tested and one positive clone was identified.

The insert (1.3 kb) was subcloned into pBluescript™ (Stratagene), and the nucleotide sequence was determined using a Sequenase™ version 2.0 DNA Sequencing kit (USB). Nucleotide and protein database search employed the BLAST™ (NCBI) and FASTA™ programs (GCG) with GenBank, EMBL, and SwissProt databases.

30 Northern Blot Analysis

Total RNAs (15 μ g) from various cells were isolated, electrophoresed through formaldehyde agarose gels and transferred to a nylon membrane (Hybond-N™, Amersham). The 483 bp *Xba* I fragment from the 5' end of the EPCR cDNA was labeled by random priming according to the manufacturer's

instructions (Multiprime™ DNA labeling system, Amersham) and used for hybridization.

Protein C and APC Binding to HUVEC

Endothelial cells in suspension bound

5 FL-APC, as monitored by flow cytometry, and demonstrated in Figure 1A. Binding was saturable and Ca^{2+} dependent, as shown by Figure 1B. Optimal binding required at least 1 mM Ca^{2+} . FL-APC was displaced from the cell surface by APC and protein
10 C equivalently, as shown by Figure 1C. The homologous Gla-domain containing proteins, protein S, factor X, and its active form, factor Xa, failed to displace bound FL-APC, suggesting that there is a specific binding site for APC on the endothelial
15 cell surface. Protein C binding was dependent on the Gla domain, since recombinant gla-domainless protein C (rGDPC) failed to displace FL-APC.

Detailed binding studies were also performed with ^{125}I -labeled APC and monolayers of
20 HUVEC, as shown by Figures 2A, 2B, 2C and 2D. The binding analysis indicated 77,000 sites per cell and a $K_d=30$ nM. This affinity is similar to that estimated from Figure 1.

Endothelial cell surface thrombomodulin
25 can interact with protein C and APC. The K_d (greater than 1 μM) (Hogg et al., (1992) J. Biol. Chem. 267, 703-706; Olsen et al., (1992) Biochemistry 31, 746-754), however, is much higher than that of the binding site described above with
30 respect to the new receptor. Furthermore, polyclonal and monoclonal antibodies against thrombomodulin that inhibit protein C activation did not inhibit the binding. Protein S also can interact with protein C and APC (Dahlbäck et al.,
35 (1992) Biochemistry 31, 12769-12777), but FL-APC binding to HUVEC was not influenced by protein S addition. Furthermore, polyclonal and monoclonal

antibodies to protein S did not inhibit the binding. These results indicate the binding site for protein C and APC on endothelium is distinct from these known molecules.

5 **Nucleotide and Predicted Protein
Structure Analysis of EPCR**

The insert was subcloned into pBluescript, and the nucleotide sequence was determined, as shown in Sequence ID No. 1. The
10 cdNA shown in Sequence ID No. 1 consists of 1302 bp, including a translation initiation ATG codon (AGGATGT, (Kozak, (1986) Cell 44, 283-292) at the 5'-end at nucleotides 25-27 of Sequence ID No. 1. A potential polyadenylation signal sequence,
15 AATAAA, (Proudfoot and Brownlee, (1976) Nature 263, 211-214) begins at nucleotide 1267 of Sequence ID No. 1, just 18 bp upstream of the poly(A) sequence.

The cdNA is predicted to code for a protein of 238 amino acids (Sequence ID No. 2),
20 which includes a 15 amino acid signal sequence (von Heijne, (1986) Nucleic Acids Res. 14, 4683-4690) at the N-terminal. Therefore, the mature protein is predicted to contain 223 amino acids. Sequence ID No. 2 is the predicted amino acid sequence of EPCR.
25 Amino acids 1-15 of Sequence ID No. 2

(MLTTLLPILLLSGWA) are the putative signal sequence determined by the method of von Heijne (von Heijne, 1986). Amino acids 211-236 of Sequence ID No. 2 (LVLGVLVGGFIIAGVAVGIFLCTGGR) are the putative
30 transmembrane domain. Potential N-glycosylation sites are present at amino acids 47-49, 64-66, 136-138, and 172-174 of Sequence ID No. 2.

Extracellular cysteine residues are present at amino acids 17, 114, 118, and 186 of Sequence ID
35 No. 2. A potential transmembrane region (Engelman et al., (1986) Annu. Rev. Biophys. Chem. 15, 321-53) consisting of 23 amino acids was identified at

the C-terminal end (beginning at amino acid 216 of Sequence ID No. 2).

The protein is predicted to be a type 1 transmembrane protein. The extracellular domain contains four potential N-glycosylation sites and four Cys residues. The cytoplasmic region contains only three amino acids and terminates with a Cys, which could be acylated to something or involved in heterodimer formation with another peptide.

Although described with reference to cloning and expression of the protein encoding sequence, larger amounts of protein can be obtained by expression in suitable recombinant host systems, such as mammalian, yeast, bacteria, or insect cells. Isolation can be facilitated by making antibodies to the recombinant protein which are then immobilized on substrates for use in purification of additional receptors, as described below.

As used herein, the nucleotide sequences encoding the receptor include the sequence shown in Sequence ID No. 1, and sequences having conservative substitutions, additions or deletions thereof which hybridize to Sequence ID No. 1 under stringent conditions. As used herein, the amino acids sequences constituting the receptor include the sequence shown in Sequence ID No. 2, and sequences having conservative substitutions, additions or deletions thereof which form a receptor having functionally equivalent biological activity. It is well known to those skilled in the art what constitutes conservative substitutions, additions or deletions, and which could be readily ascertained as encoding, or forming, a functionally equivalent receptor molecule using the functional assays described herein.

The hydropathic plot shown in Figures 4a-4c was performed according to the method of Goldman et al (Engelman et al., 1994) (solid line) and that of Kyte and Doolittle (1982) J. Mol. Biol. 157, 105-132 (dotted line).

DNA and protein database searches revealed that the sequence is related to the centrosome-associated, cell cycle dependent murine protein, CCD41, also referred as centrocyclin (Rothbarth et al., (1993) J. Cell Sci. 104, 19-30), as shown by Figures 5a-5b. The similarity in the published sequence of murine CCD41 with human EPCR led to the cloning and sequencing of the murine EPCR. The sequence of murine EPCR is shown in Figure 6. It is distinct from the published sequence of CCD41.

The EPCR amino acid sequence was also related to, but quite distinct from, the CD1/MHC superfamily and the murine CD1.2, as also shown by Figures 5a-5b. Based on the homology to CD1/MHC, it is likely that EPCR contains two domains, consisting of residues 17-114 and 118-188. Of the CD1 family members, CD1d is the most similar to EPCR. In the mouse, CCD41 is associated exclusively with the centrosome during G₁ but becomes detectable elsewhere during the cell cycle, reaching a maximum during G₂, except during the G₂/M phase (Rothbarth et al., 1993). EPCR expression appears restricted to endothelium, which would not be expected for a cell cycle associated protein.

The identification of the protein C receptor on endothelium suggests that the endothelial cell binds protein C/APC through three distinct mechanisms. In addition to EPCR, protein S can bind APC/protein C on negatively charged membrane surfaces that include the endothelium (Stern et al., (1986) J. Biol. Chem. 261, 713-718),

but this is not cell type specific (Dahlbäck et al., 1992). Thrombomodulin in complex with thrombin can bind protein C and APC (Hogg et al., 1992). On endothelium, the protein S binding sites (Nawroth and Stern, (1986) J. Exp. Med. 163, 740-745), thrombomodulin (Esmon, 1989) and EPCR are all down regulated by cytokines, indicating that inflammation can impair protein C pathway function at multiple levels.

10 The homology to the CD1/MHC family of proteins is especially interesting since it provides indications as to the function of EPCR. The CD1/MHC family has three extracellular domains termed $\alpha 1$, 2 and 3. The extracellular domain of EPCR 15 contains four Cys residues that appear to correspond to two distinct domains. EPCR lacks the third domain of the CD1/MHC family, but the two domains have significant homology to the $\alpha 1$ and $\alpha 2$ domains of the CD1 protein family and the $\alpha 2$ domain 20 of the MHC class 1 protein, suggesting that these proteins evolved from a common ancestor. The first domain of EPCR, residues 17-114, contains two potential N glycosylation sites and is rich in β strand structure, suggesting that it may form a β 25 sheet. Despite the β strand structure, consensus sequences (Williams and Barclay, (1988) Ann. Rev. Immunol. 6, 381-405) for the immunoglobulin

superfamily of receptors are absent. The second domain of EPCR, residues 118-188, contains two 30 additional N glycosylation sites and, like the CD1/MHC family, this domain is predicted to have limited β structure.

II. Modulation of Inflammation using EPCR.

In vitro studies have suggested 35 anti-inflammatory activities for APC. For instance, an unusual carbohydrate sequence on protein C can inhibit inflammatory cell adhesion to

selectins (Grinnell et al., (1994) Glycobiology, 4, 221-226). Modest inhibitory effects of APC have been reported on TNE production (Grey et al., (1993) Transplant. Proc. 25, 2913-2914). EPCR could contribute to these anti-inflammatory mechanisms. Since the homologous protein family, CD1, can be linked to CD8 (Ledbetter et al., (1985) J. Immunol. 134, 4250-4254), it is also possible that the proteins C receptor is linked to another protein and signal through this second protein. One of the CD1 family members, CD1d, has been reported to promote T cell responses, possibly involving binding to CD8 (Panja et al., (1993) J. Exp. Med. 178, 1115-1119). CD1b has recently been reported to serve as an antigen presenting molecule (Porcelli et al., (1992) Nature 360, 593-597). The ability to bind protein C/APC could then be linked either directly or indirectly to signalling via direct interaction with cells of the immune system. Since the MHC class of proteins is involved in presentation of proteins to cell receptors, the concept of presentation of protein C/APC to inflammatory cells as a means of elaborating anti-inflammatory activity may also be involved. This includes modulation of enzyme specificity such as occurs with thrombin-thrombomodulin interaction (Esmon, 1989). In this case, the EPCR-APC complex might cleave biologically active peptides from unknown substrates.

30 EPCR mRNA Levels and APC Binding

To determine the cellular specificity of EPCR expression, the intensity of FL-APC binding to HUVEC was compared to several human cell lines. Fl-APC bound strongly only to HUVEC, and not to any of the T, B, or monocytic cell lines tested. Cells were incubated at room temperature without or with 160 nM Fl-APC in the presence of 1.3 mM CaCl₂.

Binding was analyzed by flow cytometry. Slight binding was demonstrated with the osteosarcoma line, HOS and the epidermoid carcinoma cell line, HEP-2.

5 Total RNA was extracted from these cells and hybridized with the EPCR cDNA probe for Northern Blot Analysis. EPCR mRNA was detected by Northern blot analysis for HUVEC, Jurkat, HEP-2, Raji, HOS, and U937. Among the cells lines tested, 10 EPCR mRNA was detected at high levels only in HUVEC. The calculated mRNA size of 1.3 kb was identical to the size of the isolated cDNA. After prolonged exposure, a weak signal was also detected with the osteosarcoma cell line HOS and monocyte 15 cell line U937. Thus, both APC binding and EPCR mRNA expression are very specific for endothelium.

Effects of TNF on APC Binding and EPCR mRNA Levels

Several other members of the protein C 20 anticoagulant pathway are subject to regulation by inflammatory cytokines (Esmon, 1989). For instance, endothelial cell surface thrombomodulin expression and message are known to be reduced by exposure of the cells to TNF (Conway and Rosenberg, 25 1988; Lentz et al., 1991). To determine if a similar process occurs with EPCR, HUVEC were treated with TNF and APC binding and expression of EPCR mRNA were examined. APC binding to HUVEC decreased in a time dependent fashion. EPCR 30 activity decreased more rapidly than thrombomodulin antigen. HUVEC were cultured for 0, 6, 24 and 48 hr, in the presence of TNF- α (10 ng/ml). Cells were harvested and residual F1-APC binding or thrombomodulin (TM) expression was analyzed by flow 35 cytometry. Cell surface TM was stained with an anti-TM murine monoclonal antibody and FITC-conjugated anti-mouse IgG. The negative control is without added fluorescent ligand.

HUVEC were treated with 10 ng/ml of TNF- α for 0, 0.5, 1, 2, 3, 6, 10 and 24 hr, and message was extracted and detected as described above. The results demonstrated that the concentration of EPCR mRNA was also reduced by TNF treatment. Message levels and APC binding activity decreased in parallel. Therefore, the TNF mediated down-regulation of APC binding to endothelium probably occurs at the level of mRNA expression.

Enhancement of inflammatory responses by blocking binding of endogenous molecules to EPCR can be achieved by administration of compounds binding to the receptor to a subject in need of inhibition. The degree of binding is routinely determined using assays such as those described above. Compounds which are effective include antibodies to the protein, fragments of antibodies retaining the binding regions, and peptide fragments of APC which include the Gla region.

Inhibition of the inflammatory response could be obtained by infusing soluble EPCR. Alternatively, localizing EPCR to surfaces in contact with blood would render the surfaces anticoagulant by virtue of the ability of EPCR to bind and concentrate the anticoagulant APC at the surface. Alternatively, the function of EPCR could be enhanced by overexpressing the EPCR in endothelium used to coat vascular grafts in patients with vascular disease or on stents in cardiac patients.

The DNA sequence can also be used for screening for other homologous or structurally similar receptor proteins using hybridization probes. These methods and reagents and pharmaceuticals are more readily understood by reference to the following.

Screening of patient samples for expression of receptor proteins.

Patients with thrombosis or hyperinflammatory conditions could be screened for defects in the EPCR gene. Sequence ID No. 1, and consecutive portions thereof of at least about seven nucleotides, more preferably fourteen to seventeen nucleotides, most preferably about twenty nucleotides, are useful in this screening using hybridization assays of patient samples, including blood and tissues. Screening can also be accomplished using antibodies, typically labelled with a fluorescent, radiolabelled, or enzymatic label, or by isolation of target cells and screening for binding activity as described in the examples above. Typically, one would screen for expression on either a qualitative or quantitative basis, and for expression of functional receptor. Labelling can be with ^{32}P , ^{35}S , fluorescein, biotin, or other labels routinely used with methods known to those skilled in the art for labelling of proteins and/or nucleic acid sequences.

Hybridization Probes

Reaction conditions for hybridization of an oligonucleotide probe or primer to a nucleic acid sequence vary from oligonucleotide to oligonucleotide, depending on factors such as oligonucleotide length, the number of G and C nucleotides, and the composition of the buffer utilized in the hybridization reaction. Moderately stringent hybridization conditions are generally understood by those skilled in the art as conditions approximately 25°C below the melting temperature of a perfectly base-paired double-stranded DNA. Higher specificity is generally achieved by employing incubation conditions having higher temperatures, in other words, more stringent conditions. In general, the longer the sequence

or higher the G and C content, the higher the temperature and/or salt concentration required. Chapter 11 of the well-known laboratory manual of Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, second edition, Cold Spring Harbor Laboratory Press, New York (1990) (which is incorporated by reference herein), describes hybridization conditions for oligonucleotide probes and primers in great detail, including a description of the factors involved and the level of stringency necessary to guarantee hybridization with specificity. The preferred size of a hybridization probe is from 10 nucleotides to 100,000 nucleotides in length. Below 10 nucleotides, hybridized systems are not stable and will begin to denature above 20°C. Above 100,000 nucleotides, one finds that hybridization (renaturation) becomes a much slower and incomplete process, as described in greater detail in the text MOLECULAR GENETICS, Stent, G.S. and R. Calender, pp. 213-219 (1971). Ideally, the probe should be from 20 to 10,000 nucleotides. Smaller nucleotide sequences (20-100) lend themselves to production by automated organic synthetic techniques. Sequences from 100-10,000 nucleotides can be obtained from appropriate restriction endonuclease treatments. The labeling of the smaller probes with the relatively bulky chemiluminescent moieties may in some cases interfere with the hybridization process.

Generation of Antibodies for Diagnostic or Therapeutic Use

Antibodies to the receptor proteins can also be generated which are useful in detection, characterization or isolation of receptor proteins, as well as for modifying receptor protein activity, in most cases, through inhibition of binding. Antibodies are generated by standard techniques,

using human or animal receptor proteins. Since the proteins exhibit high evolutionary conservation, it may be advantageous to generate antibodies to a protein of a different species of origin than the species in which the antibodies are to be tested or utilized, looking for those antibodies which are immunoreactive with the most evolutionarily conserved regions. Antibodies are typically generated by immunization of an animal using an adjuvant such as Freund's adjuvant in combination with an immunogenic amount of the protein administered over a period of weeks in two to three week intervals, then isolated from the serum, or used to make hybridomas which express the antibodies in culture. Because the methods for immunizing animals yield antibody which is not of human origin, the antibodies could elicit an adverse effect if administered to humans. Methods for "humanizing" antibodies, or generating less immunogenic fragments of non-human antibodies, are well known. A humanized antibody is one in which only the antigen-recognized sites, or complementarily-determining hypervariable regions (CDRs) are of non-human origin, whereas all framework regions (FR) of variable domains are products of human genes. These "humanized" antibodies present a lesser xenographic rejection stimulus when introduced to a human recipient. To accomplish humanization of a selected mouse monoclonal antibody, the CDR grafting method described by Daugherty, et al., (1991) Nucl. Acids Res., 19:2471-2476, incorporated herein by reference, may be used. Briefly, the variable region DNA of a selected animal recombinant anti-idiotypic ScFv is sequenced by the method of Clackson, T., et al., (1991) Nature, 352:624-688, incorporated herein by reference. Using this

sequence, animal CDRs are distinguished from animal framework regions (FR) based on locations of the CDRs in known sequences of animal variable genes. Kabat, H.A., et al., Sequences of Proteins of Immunological Interest, 4th Ed. (U.S. Dept. Health and Human Services, Bethesda, MD, 1987). Once the animal CDRs and FR are identified, the CDRs are grafted onto human heavy chain variable region framework by the use of synthetic oligonucleotides and polymerase chain reaction (PCR) recombination. Codons for the animal heavy chain CDRs, as well as the available human heavy chain variable region framework, are built in four (each 100 bases long) oligonucleotides. Using PCR, a grafted DNA sequence of 400 bases is formed that encodes for the recombinant animal CDR/human heavy chain FR protection.

The immunogenic stimulus presented by the monoclonal antibodies so produced may be further decreased by the use of Pharmacia's (Pharmacia LKB Biotechnology, Sweden) "Recombinant Phage Antibody System" (RPAS), which generates a single-chain Fv fragment (ScFv) which incorporates the complete antigen-binding domain of the antibody. In the RPAS, antibody variable heavy and light chain genes are separately amplified from the hybridoma mRNA and cloned into an expression vector. The heavy and light chain domains are co-expressed on the same polypeptide chain after joining with a short linker DNA which codes for a flexible peptide.

This assembly generates a single-chain Fv fragment (ScFv) which incorporates the complete antigen-binding domain of the antibody. Compared to the intact monoclonal antibody, the recombinant ScFv includes a considerably lower number of epitopes, and thereby presents a much weaker immunogenic stimulus when injected into humans.

The antibodies can be formulated in standard pharmaceutical carriers for administration to patients in need thereof. These include saline, phosphate buffered saline, and other aqueous carriers, and liposomes, polymeric microspheres and other controlled release delivery devices, as are well known in the art. The antibodies can also be administered with adjuvant, such as muramyl dipeptide or other materials approved for use in humans (Freund's adjuvant can be used for administration of antibody to animals).

Screening for drugs modifying or altering the extent of receptor function or expression

The receptor proteins are useful as targets for compounds which turn on, or off, or otherwise regulate binding to these receptors. The assays described above clearly provide routine methodology by which a compound can be tested for an inhibitory effect on binding of PC or APC. The in vitro studies of compounds which appear to inhibit binding selectively to the receptors are then confirmed by animal testing. Since the molecules are so highly evolutionarily conserved, it is possible to conduct studies in laboratory animals such as mice to predict the effects in humans.

In cases where inflammatory mediators or vascular disease down regulate EPCR, it would be advantageous to increase its concentration in vivo on endothelium. The binding assays described here and the gene sequence allow assays for increased EPCR expression. Similar approaches have been taken with thrombomodulin and investigators have shown that cyclic AMP (Maruyama, I. et al. (1991) Thrombosis Research 61, 301-310) and interleukin 4 (Kapiotis, S. et al., (1991) Blood 78, 410-415) can elevate thrombomodulin expression. The ability to

screen such drugs or drugs that block TNF down regulation of EPCR provide an approach to elevating EPCR expression *in vivo* and thus enhancing anticoagulant and anti-inflammatory activity.

5 Studies based on inhibition of binding are predictive for indirect effects of alteration of receptor binding. For example, inhibition of binding of APC or increased expression of TNF is predictive of inhibition of EPCR function.

10 Assays for testing compounds for useful activity can be based solely on interaction with the receptor protein, preferably expressed on the surface of transfected cells such as those described above. Proteins in solution or
15 immobilized on inert substrates can also be utilized. These can be used to detect inhibition or enhancement in binding of PC or APC

 Alternatively, the assays can be based on interaction with the gene sequence encoding the
20 receptor protein, preferably the regulatory sequences directing expression of the receptor protein. For example, antisense which binds to the regulatory sequences, and/or to the protein encoding sequences can be synthesized using
25 standard oligonucleotide synthetic chemistry. The antisense can be stabilized for pharmaceutical use using standard methodology (encapsulation in a liposome or microsphere; introduction of modified nucleotides that are resistant to degradation or
30 groups which increase resistance to endonucleases, such as phosphorothiodates and methylation), then screened initially for alteration of receptor activity in transfected or naturally occurring cells which express the receptor, then *in vivo* in
35 laboratory animals. Typically, the antisense would inhibit expression. However, sequences which block

those sequences which "turn off" synthesis can also be targeted.

The receptor protein for study can be isolated from either naturally occurring cells or cells which have been genetically engineered to express the receptor, as described in the examples above. In the preferred embodiment, the cells would have been engineered using the intact gene.

10 *Random generation of receptor or receptor encoding sequence binding molecules.*

Molecules with a given function, catalytic or ligand-binding, can be selected for from a complex mixture of random molecules in what has been referred to as "in vitro genetics" (Szostak, (1992) TIBS 19:89). One synthesizes a large pool of molecules bearing random and defined sequences and subjects that complex mixture, for example, approximately 10^{15} individual sequences in 20 100 μ g of a 100 nucleotide RNA, to some selection and enrichment process. For example, by repeated cycles of affinity chromatography and PCR amplification of the molecules bound to the ligand on the column, Ellington and Szostak (1990) 25 estimated that 1 in 10^{10} RNA molecules folded in such a way as to bind a given ligand. DNA molecules with such ligand-binding behavior have been isolated (Ellington and Szostak, 1992; Bock et al, 1992).

30 *Computer assisted drug design*

Computer modeling technology allows visualization of the three-dimensional atomic structure of a selected molecule and the rational design of new compounds that will interact with the 35 molecule. The three-dimensional construct typically depends on data from x-ray crystallographic analyses or NMR imaging of the selected molecule. The molecular dynamics require

force field data. The computer graphics systems enable prediction of how a new compound will link to the target molecule and allow experimental manipulation of the structures of the compound and target molecule to perfect binding specificity. Prediction of what the molecule-compound interaction will be when small changes are made in one or both requires molecular mechanics software and computationally intensive computers, usually coupled with user-friendly, menu-driven interfaces between the molecular design program and the user.

Examples of molecular modelling systems are the CHARMM and QUANTA programs, Polygen Corporation, Waltham, MA. CHARMM performs the energy minimization and molecular dynamics functions. QUANTA performs the construction, graphic modelling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with each other.

A number of articles review computer modeling of drugs interactive with specific proteins, such as Rotivinen, et al., (1988) Acta Pharmaceutica Fennica 97, 159-166; Ripka, New Scientist 54-57 (June 16, 1988); McKinally and Rossmann, (1989) Annu. Rev. Pharmacol. Toxicol. 29, 111-122; Perry and Davies, QSAR: Quantitative Structure-Activity Relationships in Drug Design pp. 189-193 (Alan R. Liss, Inc. 1989); Lewis and Dean, (1989) Proc. R. Soc. Lond. 236, 125-140 and 141-162; and, with respect to a model receptor for nucleic acid components, Askew, et al., (1989) J. Am. Chem. Soc. 111, 1082-1090. Other computer programs that screen and graphically depict chemicals are available from companies such as BioDesign, Inc., Pasadena, CA., Allelix, Inc., Mississauga, Ontario, Canada, and Hypercube, Inc.,

Cambridge, Ontario. Although these are primarily designed for application to drugs specific to particular proteins, they can be adapted to design of drugs specific to regions of DNA or RNA, once that region is identified.

Although described above with reference to design and generation of compounds which could alter binding, one could also screen libraries of known compounds, including natural products or synthetic chemicals, and biologically active materials, including proteins, for compounds which are inhibitors or activators.

Generation of nucleic acid regulators

Nucleic acid molecules containing the 5' regulatory sequences of the receptor genes can be used to regulate or inhibit gene expression in vivo. Vectors, including both plasmid and eukaryotic viral vectors, may be used to express a particular recombinant 5' flanking region-gene construct in cells depending on the preference and judgment of the skilled practitioner (see, e.g., Sambrook et al., Chapter 16). Furthermore, a number of viral and nonviral vectors are being developed that enable the introduction of nucleic acid sequences in vivo (see, e.g., Mulligan, (1993) Science, 260, 926-932; United States Patent No. 4,980,286; United States Patent No. 4,868,116; incorporated herein by reference). Recently, a delivery system was developed in which nucleic acid is encapsulated in cationic liposomes which can be injected intravenously into a mammal. This system has been used to introduce DNA into the cells of multiple tissues of adult mice, including endothelium and bone marrow (see, e.g., Zhu et al., (1993) Science 261, 209-211; incorporated herein by reference).

The 5' flanking sequences of the receptor gene can also be used to inhibit the expression of the receptor. For example, an antisense RNA of all or a portion of the 5' flanking region of the receptor gene can be used to inhibit expression of the receptor *in vivo*. Expression vectors (e.g., retroviral expression vectors) are already available in the art which can be used to generate an antisense RNA of a selected DNA sequence which is expressed in a cell (see, e.g., U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286). Accordingly, DNA containing all or a portion of the sequence of the 5' flanking region of the receptor gene can be inserted into an appropriate expression vector so that upon passage into the cell, the transcription of the inserted DNA yields an antisense RNA that is complementary to the mRNA transcript of the receptor protein gene normally found in the cell. This antisense RNA transcript of the inserted DNA can then base-pair with the normal mRNA transcript found in the cell and thereby prevent the mRNA from being translated. It is of course necessary to select sequences of the 5' flanking region that are downstream from the transcriptional start sites for the receptor protein gene to ensure that the antisense RNA contains complementary sequences present on the mRNA. Antisense RNA can be generated *in vitro* also, and then inserted into cells. Oligonucleotides can be synthesized on an automated synthesizer (e.g., Model 8700 automated synthesizer of Milligen-Bioscience, Burlington, MA or ABI Model 380B). In addition, antisense deoxyoligonucleotides have been shown to be effective in inhibiting gene transcription and viral replication (see e.g., Zamecnik et al.,

(1978) Proc. Natl. Acad. Sci. USA 75, 280-284;
Zamecnik et al., (1986) Proc. Natl. Acad. Sci., 83,
4143-4146; Wickstrom et al., (1988) Proc. Natl.
Acad. Sci. USA 85, 1028-1032; Crooke, (1993) FASEB
J. 7, 533-539. Furthermore, recent work has shown
that improved inhibition of expression of a gene by
antisense oligonucleotides is possible if the
antisense oligonucleotides contain modified
nucleotides (see, e.g., Offensperger et al.,
(1993) EMBO J. 12, 1257-1262 (in vivo inhibition of
duck hepatitis B viral replication and gene
expression by antisense phosphorothioate
oligodeoxynucleotides); PCT WO 93/01286 Rosenberg
et al., (synthesis of sulfurthioated
oligonucleotides); Agrawal et al., (1988) Proc.
Natl. Acad. Sci. USA 85, 7079-7083 (synthesis of
antisense oligonucleoside phosphoramidates and
phosphorothioates to inhibit replication of human
immunodeficiency virus-1); Sarin et al., (1989)
Proc. Natl. Acad. Sci. USA 85, 7448-7794 (synthesis
of antisense methylphosphonate oligonucleotides);
Shaw et al., (1991) Nucleic Acids Res 19, 747-750
(synthesis of 3' exonuclease-resistant
oligonucleotides containing 3' terminal
phosphoroamidate modifications); incorporated
herein by reference). The sequences of the 5' flanking region
of receptor protein gene can also be used in triple
helix (triplex) gene therapy. Oligonucleotides
complementary to gene promoter sequences on one of
the strands of the DNA have been shown to bind
promoter and regulatory sequences to form local
triple nucleic acid helices which block
transcription of the gene (see, e.g., Maher et al.,
(1989) Science 245, 725-730; Orson et al., (1991)
Nucl. Acids Res. 19, 3435-3441; Postal et al.,
(1991) Proc. Natl. Acad. Sci. USA 88, 8227-8231;

Cooney et al., (1988) Science 241, 456-459; Young et al., (1991) Proc. Natl. Acad. Sci. USA 88, 10023-10026; Duval-Valentin et al., (1992) Proc. Natl. Acad. Sci. USA 89, 504-508; Blume et al.,
5 (1992) Nucl. Acids Res. 20, 1777-1784; Grigoriev et al., (1992) J. Biol. Chem. 267, 3389-3395.

Recently, both theoretical calculations and empirical findings have been reported which provide guidance for the design of oligonucleotides
10 for use in oligonucleotide-directed triple helix formation to inhibit gene expression. For example, oligonucleotides should generally be greater than 14 nucleotides in length to ensure target sequence specificity (see, e.g., Maher et al., (1989);
15 Grigoriev et al., (1992)). Also, many cells avidly take up oligonucleotides that are less than 50 nucleotides in length (see e.g., Orson et al., (1991); Holt et al., (1988) Mol. Cell. Biol. 8, 963-973; Wickstrom et al., (1988) Proc. Natl. Acad. Sci. USA 85, 1028-1032). To reduce susceptibility
20 to intracellular degradation, for example by 3' exonucleases, a free amine can be introduced to a 3' terminal hydroxyl group of oligonucleotides without loss of sequence-binding specificity (Orson
25 et al., 1991). Furthermore, more stable triplexes are formed if any cytosines that may be present in the oligonucleotide are methylated, and also if an intercalating agent, such as an acridine
30 derivative, is covalently attached to a 5' terminal phosphate (e.g., via a pentamethylene bridge); again without loss of sequence specificity (Maher et al., (1989); Grigoriev et al., (1992)).

Methods to produce or synthesize oligonucleotides are well known in the art. Such
35 methods can range from standard enzymatic digestion followed by nucleotide fragment isolation (see e.g., Sambrook et al., Chapters 5, 6) to purely

synthetic methods, for example, by the cyanoethyl phosphoramidite method using a Milligen or Beckman System 1Plus DNA synthesizer (see also, Ikuta et al., (1984) Ann. Rev. Biochem. 53, 323-356 (phosphotriester and phosphite-triester methods); Narang et al., (1980) Methods Enzymol., 65, 610-620 (phosphotriester method). Accordingly, DNA sequences of the 5' flanking region of the receptor protein gene described herein can be used to design and construct oligonucleotides including a DNA sequence consisting essentially of at least 15 consecutive nucleotides, with or without base modifications or intercalating agent derivatives, for use in forming triple helices specifically within the 5' flanking region of a receptor protein gene in order to inhibit expression of the gene.

In some cases it may be advantageous to insert enhancers or multiple copies of the regulatory sequences into an expression system to facilitate screening of methods and reagents for manipulation of expression.

Preparation of Receptor Protein Fragments

Compounds which are effective for blocking binding of the receptor can also consist of fragments of the receptor proteins, expressed recombinantly and cleaved by enzymatic digest or expressed from a sequence encoding a peptide of less than the full length receptor protein. These will typically be soluble proteins, i.e., not including the transmembrane and cytoplasmic regions, although smaller portions determined in the assays described above to inhibit or compete for binding to the receptor proteins can also be utilized. It is a routine matter to make appropriate receptor protein fragments, test for binding, and then utilize. The preferred fragments are of human origin, in order to minimize potential

immunological response. The peptides can be as short as five to eight amino acids in length and are easily prepared by standard techniques. They can also be modified to increase in vivo half-life, by chemical modification of the amino acids or by attachment to a carrier molecule or inert substrate. Based on studies with other peptide fragments blocking receptor binding, the IC_{50} , the dose of peptide required to inhibit binding by 50%, ranges from about 1 μM to greater than 10 mM, depending on the peptide size and folding. These ranges are well within the effective concentrations for the in vivo administration of peptides, based on comparison with the RGD-containing peptides, described, for example, in U.S. Patent No. 4,792,525 to Ruoslahti, et al., used in vivo to alter cell attachment and phagocytosis. The peptides can also be conjugated to a carrier protein such as keyhole limpet hemocyanin by its N-terminal cysteine by standard procedures such as the commercial Imject kit from Pierce Chemicals or expressed as a fusion protein, which may have increased efficacy.

As noted above, the peptides can be prepared by proteolytic cleavage of the receptor proteins, or, preferably, by synthetic means. These methods are known to those skilled in the art. An example is the solid phase synthesis described by J. Merrifield, (1964) J. Am. Chem. Soc. 85, 2149, used in U.S. Patent No. 4,792,525, and described in U.S. Patent No. 4,244,946, wherein a protected alpha-amino acid is coupled to a suitable resin, to initiate synthesis of a peptide starting from the C-terminus of the peptide. Other methods of synthesis are described in U.S. Patent No. 4,305,872 and 4,316,891. These methods can be used to synthesize peptides having identical

sequence to the receptor proteins described herein, or substitutions or additions of amino acids, which can be screened for activity as described above.

The peptide can also be administered as a
5 pharmaceutically acceptable acid- or base- addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as
10 formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide,
15 potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

Peptides containing cyclopropyl amino acids, or amino acids derivatized in a similar
20 fashion, can also be used. These peptides retain their original activity but have increased half-lives in vivo. Methods known for modifying amino acids, and their use, are known to those skilled in the art, for example, as described in U.S. Patent
25 No. 4,629,784 to Stammer.

The peptides are generally active when administered parenterally in amounts above about 1 $\mu\text{g/kg}$ of body weight. Based on extrapolation from other proteins, for treatment of most inflammatory
30 disorders, the dosage range will be between 0.1 to 70 mg/kg of body weight. This dosage will be dependent, in part, on whether one or more peptides are administered.

Pharmaceutical Compositions

35 Compounds which alter receptor protein binding are preferably administered in a pharmaceutically acceptable vehicle. Suitable

pharmaceutical vehicles are known to those skilled in the art. For parenteral administration, the compound will usually be dissolved or suspended in sterile water or saline. For enteral

5 administration, the compound will be incorporated into an inert carrier in tablet, liquid, or capsular form. Suitable carriers may be starches or sugars and include lubricants, flavorings, binders, and other materials of the same nature.

10 The compounds can also be administered locally by topical application of a solution, cream, gel, or polymeric material (for example, a Pluronic™, BASF).

Alternatively, the compound may be
15 administered in liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a patient are known to those skilled in the art. U.S. Patent No. 4,789,734 describe methods for encapsulating
20 biological materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed or sonicated, as necessary. A
25 review of known methods is by G. Gregoriadis, Chapter 14. "Liposomes", Drug Carriers in Biology and Medicine pp. 287-341 (Academic Press, 1979). Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be
30 tailored for passage through the gastrointestinal tract directly into the bloodstream.

Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time,
35 ranging from days to months. See, for example, U.S. Patent Nos. 4,906,474, 4,925,673, and 3,625,214.

~~Disorders to be treated~~
As described herein, a variety of compounds can be used to inhibit or enhance expression of the EPCR. The nature of the disorder will determine if the expression should be enhanced or inhibited. For example, based on the studies involving the use of an anti-protein C antibody in combination with cytokine, it should be possible to treat solid tumors by enhancing an inflammatory response involving blocking of protein C or activated protein C binding to an endothelial cell protein C/activated protein C receptor by administering to a patient in need of treatment thereof an amount of a compound blocking binding of protein C or activated protein C to the receptor. Similarly, it should be possible to treat disorders such as gram negative sepsis, stroke, thrombosis, septic shock, adult respiratory distress syndrome, and pulmonary emboli using a method for inhibiting an inflammatory response involving administration of EPCR or EPCR fragments or substances that upregulate EPCR expression to a patient in need of treatment thereof.

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Oklahoma Medical Research Foundation
 - (ii) TITLE OF INVENTION: Cloning and Regulation of an Endothelial Cell Protein C/Activated Protein C Receptor
 - (iii) NUMBER OF SEQUENCES: 2
 - (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Patrea L. Pabst
 - (B) STREET: 2800 One Atlantic Center
 - (C) CITY: Atlanta
 - (D) STATE: Georgia
 - (E) COUNTRY: USA
 - (F) ZIP: 30309-3450
 - (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
 - (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Pabst, Patrea L.
 - (B) REGISTRATION NUMBER: 31,284
 - (C) REFERENCE/DOCKET NUMBER: OMRF152
 - (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (404) 873-8794
 - (B) TELEFAX: (404) 873-8795
- (2) INFORMATION FOR SEQ ID NO:1:
- (1) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1302 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: cDNA
 - (iii) HYPOTHETICAL: NO
 - (iv) ANTI-SENSE: NO
 - (ix) FEATURE:
 - (A) NAME/KEY: misc_feature

(B) LOCATION: 1..1302

(D) OTHER INFORMATION: /note= "Nucleotides 25 through 738 encode the Endothelial Cell Protein Receptor of Sequence ID No. 2."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGGTCGGGA	GCCTCAACTT	CAGGATGTTG	ACAACATTGC	TGCCGATACT	GCTGCTGTCT	60
GGCTGGGCCT	TTTGTAGCCA	AGACGCCCTCA	GATGGCCTCC	AAAGACTTCA	TATGCTCCAG	120
ATCTCCTACT	TCCGGGACCC	CTATCACGTG	TGGTACCAGG	GCAACGCGTC	GCTGGGGGGA	180
CACCTAACGC	ACGTGCTGGA	AGGCCCAGAC	ACCAACACCA	CGATCATTCA	GCTGCAGCCC	240
TTGCAGGAGC	CCGAGAGCTG	GGCGCGCAGC	CAGAGTGGCC	TGCAGTCCTA	CCTGCTCCAG	300
TTCCACGGCC	TCGTGCGCCT	GGTGCACCAG	GAGCGGACCT	TGGCCTTTCC	TCTGACCATC	360
CGCTGCTTCC	TGGGCTGTGA	GCTGCCCTCC	GAGGGCTCTA	GAGCCCATGT	CTTCTTCGAA	420
GTGGCTGTGA	ATGGGAGCTC	CTTTGTGAGT	TTCCGGGCCGG	AGAGAGCCTT	GTGGCAGGCA	480
GACACCCAGG	TCACCTCCGG	AGTGGTCACC	TTACCCCTGC	AGCAGCTCAA	TGCCTACAAC	540
CGCACTCGGT	ATGAAC TGCG	GGAATTCCTG	GAGGACACCT	GTGTGCAGTA	TGTGCAGAAA	600
CATATTTCGG	CGGAAACAC	GAAAGGGAGC	CAAACAAGCC	GCTCCTACAC	TTGCTGGTTC	660
CTGGGCGTCC	TGGTGGGCGG	TTTCATCATT	GCTGGTGTGG	CTGTAGGCAT	CTTCCTGTGC	720
ACAGGTGGAC	GGCGATGTTA	ATTACTCTCC	AGCCCCGTCA	GAAGGGGCTG	GATTGATGGA	780
GGCTGGCAAG	GGAAAAGTTT	AGCTCACTGT	GAAAGCCAGAC	TCCCCAACTG	AAACACCAGA	840
AGGTTTGAG	TGACAGCTCC	TTTCTTCTCC	CACATCTGCC	CACIGAAGAT	TTGAGGGAGG	900
GGAGATGGAG	AGGAGAGGTG	GACAAAGTAC	TTGGTTTGCT	AAGAACCTAA	GAACGTGTAT	960

GCTTTGCTGA ATTAGTCTGA TAAGTGAATG TTTATCTATC TTTGTGGAAA ACAGATAATG 1020
 GAGTTGGGC AGGAAGCCTA TGGCCATCC TCCAAAGACA GACAGAATCA CCTGAGGCGT 1080
 TCAAAAGATA TAACCAAATA AACAAAGTCAT CCACAATCAA AATACAACAT TCAATACTTC 1140
 CAGGTGTGTC AGACTTGGGA TGGGACGCTG ATATAATAGG GTAGAAAGAA GTAACACGAA 1200
 GAAGTGGTGG AAATGTAAA TCCAAGTCAT ATGGCAGTGA TCAATTATTA ATCAATTAAAT 1260
 AATATTAAATA AATTCTTAT ATTAAAAA AAAAAAAAAA AA 1302

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 238 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

(A) NAME/KEY: misc feature

(B) LOCATION: 1..365

(D) OTHER INFORMATION: /note= "Endothelial Cell Protein Receptor encoded by

nucleotides 1 through 1302 of Sequence ID No. 1."

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 1..15

(D) OTHER INFORMATION: /note= "Amino acids 1-15 represent a putative signal sequence."

(ix) FEATURE:

(A) NAME/KEY: Domain

(B) LOCATION: 211..236

(D) OTHER INFORMATION: /note= "Amino acids 211-236 represent a putative transmembrane domain."

(ix) FEATURE:

(A) NAME/KEY: Activ -site

(B) LOCATION: 47..174

(D) OTHER INFORMATION: /note= "Amino acids 47-49, 64-66, 136-138 and 172-174 represent potential N-glycosylation sites."

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 17..186

(D) OTHER INFORMATION: /note= "Amino acids 17, 114, 118 and 186 represent extracellular cysteine residues."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Leu Thr Thr Leu Leu Pro Ile Leu Leu Leu Ser Gly Trp Ala Phe
1      5      10      15

Cys Ser Gln Asp Ala Ser Asp Gly Leu Gln Arg Leu His Met Leu Gln
20      25      30

Ile Ser Tyr Phe Arg Asp Pro Tyr His Val Trp Tyr Gln Gly Asn Ala
35      40      45

Ser Leu Gly Gly His Leu Thr His Val Leu Leu Glu Gly Pro Asp Thr Asn
50      55      60

Thr Thr Ile Ile Gln Leu Leu Gln Pro Leu Gln Glu Pro Glu Ser Trp Ala
65      70      75      80

Arg Thr Gln Ser Gly Leu Leu Gln Ser Tyr Leu Leu Gln Phe His Gly Leu
85      90      95

Val Arg Leu Val His Gln Glu Arg Thr Leu Ala Phe Pro Leu Thr Ile
100      105      110

Arg Cys Phe Leu Gly Cys Glu Leu Pro Pro Glu Gly Ser Arg Ala His
115      120      125

Val Phe Phe Glu Val Ala Val Asn Gly Ser Ser Phe Val Ser Phe Arg
130      135      140

```

Pro Glu Arg Ala Leu Trp Gln Ala Asp Thr Gln Val Thr Ser Gly Val
 145 150 155 160
 Val Thr Phe Thr Leu Gln Gln Leu Asn Ala Tyr Asn Arg Thr Arg Tyr
 165 170 175
 Glu Leu Arg Glu Phe Leu Glu Asp Thr Cys Val Gln Tyr Val Gln Lys
 180 185 190
 His Ile Ser Ala Glu Asn Thr Lys Gly Ser Gln Thr Ser Arg Ser Tyr
 195 200 205
 Thr Ser Leu Val Leu Gly Val Leu Val Gly Gly Phe Ile Ile Ala Gly
 210 215 220
 Val Ala Val Gly Ile Phe Leu Cys Thr Gly Gly Arg Arg Cys
 225 230 235

We claim.

1. An isolated endothelial cell protein C/activated protein C receptor.

2. The receptor of claim 1 encoded by the nucleotide sequence of Sequence ID No. 1 and degenerative sequences thereof and sequence having conservative substitutions, additions or deletions thereof hybridizing to Sequence ID No. 1 under stringent conditions, which encode the receptor.

3. The receptor of claim 1 having the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

4. The receptor of claim 1 expressed on the surface of a non-human cell or a non-endothelial cell.

5. The receptor of claim 1 in soluble form.

6. The receptor of claim 5 lacking at least a portion of the transmembrane region.

7. A nucleotide sequence encoding an endothelial cell protein C/activated protein C receptor.

8. The sequence of claim 7 having the nucleotide sequence of Sequence ID No. 1 or degenerative sequences thereof and sequence having conservative substitutions, additions or deletions thereof and hybridizing under stringent conditions to Sequence ID No. 1.

9. The sequence of claim 7 encoding the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

10. The sequence of claim 7 further comprising an expression vector.

11. The sequence of claim 10 further comprising an expression host.

12. The sequence of claim 10 expressed on the surface of a non-human cell or a non-endothelial cell.

13. The sequence of claim 7 encoding a soluble form of the receptor.

14. The sequence of claim 7 encoding a fragment of the receptor of at least fourteen consecutive nucleotides in length.

15. The sequence of claim 14 labelled with a detectable label.

16. A method for enhancing an inflammatory response involving blocking of protein C or activated protein C binding to an endothelial cell protein C/activated protein C receptor comprising administering to a patient in need of treatment thereof an amount of a compound blocking binding of protein C or activated protein C to the receptor.

17. The method of claim 16 wherein the compound is selected from the group consisting of antibodies and fragments of antibodies to the receptor, nucleic acid sequences inhibiting expression of the receptor, and synthetic or natural compounds other than proteins, peptides or nucleic acid sequences which inhibit binding.

18. A method for inhibiting an inflammatory response involving administration of a compound selected from the group consisting of EPCR or EPCR fragments and substances that upregulate EPCR expression to a patient in need of treatment thereof.

19. An antibody or antibody fragment specifically immunoreactive with a unique epitope of an isolated endothelial cell protein C/activated protein C receptor.

20. The antibody of claim 19 wherein the receptor is encoded by the nucleotide sequence of



Sequence ID No. 1 and degenerative sequences thereof and sequence having conservative substitutions, additions or deletions thereof and hybridizing to Sequence ID No. 1 under stringent conditions.

21. The antibody of claim 19 wherein the receptor has the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

22. A method for screening for a compound which alters the binding of an endothelial receptor protein to protein C or activated protein C comprising providing an assay for binding of protein C or activated protein C to the receptor protein, adding the compound to be tested to the assay, and

determining if the amount of protein C or activated protein C which is bound to the receptor protein is altered as compared to binding in the absence of the compound to be tested.

23. A method for screening patients for abnormal receptor protein activity or function comprising determining the presence of an endothelial cell surface receptor binding protein C and activated protein C, and comparing the receptor to determine if the quantity present or the function of the receptor is equivalent to that present in normal cells.



2/10




FIG. 2a




FIG. 2b

SUBSTITUTE SHEET (RULE 26)




FIG. 2c




FIG. 2d

SUBSTITUTE SHEET (RULE 26)

4/10




FIG. 3a




FIG. 3b

5/10

FIG. 4a

CAGGTCCGGAGCCTCAACTTCAGGATGTTGACAACATTGCTGCCGATACTGCTGCTGCT 60
M L T T L L P I L L L S 12

GGCTGGGCGCTTTGTAGCCAAGAGCGCTCAGATGGCCTCCAAGACTTCATATGCTCCAG 120
G W A F © S Q D A S D G L Q R L H M L Q 32

ATCTCCTACTTCGGGACCCCTATCAGGTGTGGTACCAGGGCAACGGCTCGCTGGGGGA 180
I S Y F R D P Y H V W Y Q G N A S L G G 52

CACCTAAGCACGTGCTGGAAGGCCAGACACCAACACGATCATTGAGTGCAGCCC 240
H L T H V L E G P D T N T T I I Q L Q P 72

TTGCAGGAGCCGAGAGCTGGGGCGCACGAGAGTGGCTGCAGTCTACCTGCTCCAG 300
L Q E P E S W A R T Q S G L Q S Y L L Q 92

TTCCACGGCCTGCTGGCGCTGGTGCACCAGGAGCGGACCTTGGCCTTCTCTGACCATC 360
F H G L V R L V H Q E R T L A F P L T I 112

CGCTGCTTCCTGGGCTGTGAGCTGCCCTCCGAGGGCTCTAGAGCCCATGCTCTTCTCGAA 420
R © F L G © E L P P E G S R A H V F F E 132

SUBSTITUTE SHEET (RULE 26)

6/10

FIG. 4b

GTGGCTGTAATGGAGCTCCTTTGTGAGTTCCGGCCGGAGAGAGCCTTGTGGCAGGCA 480
 V A V N G S S F V S F R P E R A L W Q A 152

GACACCAGGTCACCTCCGGAGTGGTCACCTTCACCTGCAGCAGCTCAATGCCTACAAC 540
 D T Q V T S G V V T F T L Q Q L N A Y N 172

CGCACTCGGTATGAACCTCGGGAATTCCTGGAGGACACCTGTGTGCAGTATGTCAGAAA 600
R T R Y E L R E F L E D T © V Q Y V Q K 192

CATATTCGGGAAACACGAAGGGAGCCAAACAGCCGCTCCTACACTTCGCTGGTC 660
 H I S A E N T K G S Q T S R S Y T S L V == 212

CTGGCGTCTGCTGGCGGTTTCATCATTTGCTGGTGGCTGTAGGCACTTCCTGTGC 720
L G V L V G G F I I A G V A V G I F L C 232


ACAGGTGACGGGATGTTAATTACTCTCAGCCCGCTCAGAAGGGGCTGGATTGATGGA 780
T G G R R C * 238

SUBSTITUTE SHEET (RULE 26)

7/10

FIG. 4c

GGCTGGCAAGGAAAGTTTCAGCTCACTGTGAAGCCAGACTCCCAACTGAAACACCAGA 840
 AGGTTGGAGTGACAGCTCCTTTCTTCCACATCTGCCACTGAAGATTGAGGGAGG 900
 GGAGATGGAGAGGAGAGGTGGACAAAGTACTTGGTTTGCTAAGAACCCTAAGAACGTGTAT 960
 GCTTTGCTGAATTAGTCTGATAAGTGAATGTTTATCTATCTTTGTGGAAACAGATAATG 1020
 GAGTTGGGCAGGAAGCCTATGGCCATCTCTCCAAAGACAGACAGAATCACCTGAGGCGT 1080
 TCAAAGATATAACCAATAAACAAGTCATCCACAATCAAAATACACATTCAATCTTC 1140
 CAGGTGTCAGACTTGGGATGGGACGCTGATAATAATAGGCTAGAAAGAAGTAACACGAA 1200
 GAAGTGGTGGAATGTAAATCCAAGTCATATGGCAGTGATCAATTATTATCAATTAAT 1260
 AATATTAATAAATTTCTTATATTAAAAAATAAAAAA 1302



8/10

FIG. 5a

```

*      *      *      *      *      *      *      *      *      *      *
**     **     **     **     **     **     **     **     **     **     **
EPCR  MLTT..LLPI..LLSGWAFCSQDASDGQRQHMLQISYFRDPYHVWYQGNASLGHLTHVL
CCD41 MLTKFLLLLULLLPGLAFVTPMAPKAAAYAPDLL...FPRPPSCEASGQRSGLKLLTHTL
CD1d  MGC...LFLLLWALLQANGSAEVPQRFPRLCLQISSFANSSWTRTDGLAWLGELQTHSW
MCD1.2 MRY..LPCLLLWAFLLQVWGQSEVQKNYTFRCLQTSFFANISWSTRDLSILLGDLOTHR

```

```

*      *      *      *      *      *      *      *      *      *      *
***   ***   ***   ***   ***   ***   ***   ***   ***   ***   ***
EPCR  EGPDTNTTIIQLQPLQEPESWARTOSGLQSYLLQFHGLVRLVHDER.....TLAFPLTI
CCD41 EGPSQNVTTIQLQPMQDPESWERTESGLQIYLTQFESLVKLVYRER.....KENVFFPLTV
CD1d  SN..DSQTVRSILRPWSQGTFSQQWETLOHIFRVYRSSFTRDVKEFKML..RLSYPLJEL
MCD1.2 SN..DSATISFTKPPWSQGLSNQQWEKLOHMFQVYRVSFTRDIQELVKMMSPKEDYPIEI

```

```

***   ***   ***   ***   ***   ***   ***   ***   ***   ***   ***
****  *      *      *      *      *      *      *      *      *      *
EPCR  RCFLGCEL..P...PEGSRAHVFFFEVAVNGSSFVSFRPERALWQADTQVTSGVMTFTLQQL
CCD41 SCSLGCEL..PEEEEGSEPHVFFDVAVNGSAFVSFRPKTAMVSGSQEPSKAANFTLRQL
CD1d  QVSAGCEVHP.....GNASNNFFHVAFQGKDIUSFQGTS..WEP.TQEAPLWNLAIQVL
MCD1.2 QLSTGCEMYPI.....GNASESFFHVAFQGKYAVRFRGTS..WQR.VLGAPSWLDLPKVL

```

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/US 95/09636

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 A61K39/395 C12N15/11 A61K38/17
C07K16/28 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages.	Relevant to claim No.
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 42, 21 October 1994 BALTIMORE, MD, USA, pages 26486-26491, K. FUKUDOME ET AL. 'Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor.' see the whole document --- -/--	1-3,7-11

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

2 January 1996

Date of mailing of the international search report

26.01.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 95/09636

C.(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	CIRCULATION; vol. 90, no. 4 part 2, October 1994 NEW YORK, NY, USA, page I-N K. FUKUDOME ET AL. 'Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor.' see the whole document	1-3,7-11
P,X	CIRCULATION, vol. 90, no. 4 part 2, October 1994 NEW YORK, NY, USA, page I133 K. FUKUDOME ET AL. 'Identification, cloning and regulation of a novel endothelial cell protein C/activated protein C receptor.' see abstract 0707	1-3,7-11
P,X	THROMBOSIS AND HAEMOSTASIS, vol. 72, no. 3, September 1994 STUTTGART, GERMANY, pages 465-474, N. BANGALORE ET AL. 'High affinity binding sites for activated protein C and protein C on cultured human umbilical vein endothelial cells.' see summary see page 472, right column, line 9 - line 50	1-3,7-11
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 10, 10 March 1995 BALTIMORE, MD, USA, pages 5571-5577, K. FUKUDOME ET AL. 'Molecular cloning and expression of murine and bovine endothelial cell protein C/activated protein C receptor (EPCR).' see abstract see figures 1-5	1-4,7-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/09636

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-18
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although these claims are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a):

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**CORRECTED
VERSION**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/12, C07K 14/705, A61K 39/395, C12N 15/11, A61K 38/17, C07K 16/28, G01N 33/68	A1	(11) International Publication Number: WO 96/05303 (43) International Publication Date: 22 February 1996 (22.02.96)
(21) International Application Number: PCT/US95/09636 (22) International Filing Date: 9 August 1995 (09.08.95) (30) Priority Data: 289,699 12 August 1994 (12.08.94) US (71) Applicant: OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 N.E. Thirteenth Street, Oklahoma City, OK 73104 (US). (72) Inventors: FUKUDOME, Kenji; 13125 West Park Place, Oklahoma City, OK 73142 (US). ESMON, Charles, T.; 5800 North Stonewall, Oklahoma City, OK 73111 (US). (74) Agent: PABST, Patrea, L.; Amall Golden & Gregory, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).	(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: CLONING AND REGULATION OF AN ENDOTHELIAL CELL PROTEIN C/ACTIVATED PROTEIN C RECEPTOR

(57) Abstract

Human protein C and activated protein C were shown to bind to endothelium specifically, selectively and saturably ($K_d=30$ nM, 7000 sites per cell) in a Ca^{2+} dependent fashion. Expression cloning revealed a 1.3 kb cDNA that coded for a novel type I transmembrane glycoprotein capable of binding protein C. This protein appears to be a member of the CD1/MHC superfamily. Like thrombomodulin, the receptor involved in protein C activation, the endothelial cell protein C receptor (EPCR) function and message are both down regulated by exposure of endothelium to TNF. Identification of EPCR as a member of the CD1/MHC superfamily provides insights into the role of protein C in regulating the inflammatory response, and determination of methods for pharmaceutical use in manipulating the inflammatory response.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

CLONING AND REGULATION OF AN ENDOTHELIAL CELL
PROTEIN C/ACTIVATED PROTEIN C RECEPTOR

Background of the Invention

The present invention is generally in the area of cloning, expression, and regulation of an endothelial cell protein C/activated protein C
5 receptor.

Protein C plays a major role in the regulation of blood coagulation. Patients deficient in protein C usually exhibit life threatening thrombotic-complications in infancy
10 (Seligsohn et al., (1984) N. Engl. J. Med. 310, 559-562; Esmon, (1992) Trends Cardiovasc. Med. 2, 214-220) that are corrected by protein C administration (Dreyfus et al., (1991) N. Engl. J. Med. 325, 1565-1568). In addition, activated
15 protein C (APC) can prevent the lethal effects of *E. coli* in baboon models of gram negative sepsis (Taylor et al., (1987) J. Clin. Invest. 79; U.S. Patent No. 5,009,889 to Taylor and Esmon) and preliminary clinical results suggest that protein C
20 is effective in treating certain forms of human septic shock (Gerson et al., (1993) Pediatrics 91, 418-422). These results suggest that protein C may both control coagulation and influence inflammation. Indeed, inhibition of protein S, an
25 important component of the protein C pathway, exacerbates the response of primates to sublethal levels of *E. coli* and augments the appearance of TNF in the circulation (Taylor et al., (1991) Blood 78, 357-363). The mechanisms involved in
30 controlling the inflammatory response remain unknown.

Protein C is activated when thrombin, the terminal enzyme of the coagulation system, binds to an endothelial cell surface protein, thrombomodulin
35 (Esmon, (1989) J. Biol. Chem. 264, 4743-4746; Dittman and Majerus, (1990) Blood 75, 329-336;

Dittman, (1991) Trends Cardiovasc. Med. 1, 331-336). In cell culture, thrombomodulin transcription is blocked by exposure of endothelial cells to tumor necrosis factor (TNF) (Conway and Rosenberg, (1988) Mol. Cell. Biol. 8, 5588-5592) and thrombomodulin activity and antigen are subsequently internalized and degraded (Lentz et al., (1991) Blood 77, 543-550, Moore, K.L., et al., (1989) Blood 73, 159-165). In addition, C4bBP, a regulatory protein of the complement system, binds protein S to form a complex that is functionally inactive in supporting APC anticoagulant activity in vitro (Dahlbäck, (1986) J. Biol. Chem. 261, 12022-12027) and in vivo (Taylor, et al., 1991). Furthermore, C4bBP behaves as an acute phase reactant (Dahlbäck, (1991) Thromb. Haemostas. 66, 49-61). Thus, proteins of this pathway not only appear to regulate inflammation, but they also interact with components that regulate inflammation, and they themselves are subject to down regulation by inflammatory mediators.

Given the central role of the protein C pathway in regulating the host response to inflammation and the critical role of the pathway in controlling blood coagulation, it is important to identify and characterize all of the components that interact with the system. This is especially true since the molecular basis of the anti-inflammatory effects of the protein C pathway components have yet to be elucidated at the molecular level.

It is therefore an object of the present invention to provide a cellular receptor for protein C and activated protein C.

It is a further object of the present invention to provide nucleotide sequences encoding

the cellular receptor and amino acid characterization of the receptor which allows expression of recombinant native and modified forms of the receptor.

5 It is another object of the present invention to provide methods of modulating the inflammatory response involving protein C and activated protein C.

Summary of the Invention

10 An endothelial cell protein C binding protein (referred to herein as "EPCR") has been cloned and characterized. The protein is predicted to consist of 238 amino acids, which includes a 15 amino acid signal sequence at the N-terminus, and a 15 23 amino acid transmembrane region which characterizes the receptor as a type 1 transmembrane protein. The protein binds with high affinity to both protein C and activated protein C (Kd=30 nM) and is calcium dependent. The message 20 and binding function of the receptor are both down regulated by cytokines such as TNF.

These results identify a new member of a complex pathway that, like other members of the pathway, is subject to regulation by inflammatory 25 cytokines, and can therefore be used to modulate inflammatory reactions in which protein C or activated protein C is involved. Inhibition of the inflammatory response can be obtained by infusing soluble EPCR. Alternatively, localizing 30 EPCR to surfaces in contact with blood will render the surfaces anticoagulant by virtue of the ability of EPCR to bind and concentrate the anticoagulant activated protein C at the surface. Alternatively, the function of EPCR can be enhanced by 35 overexpressing the EPCR in endothelium that could

be used to coat vascular grafts in patients with vascular disease or on stents in cardiac patients.

Brief Description of the Drawings

Figures 1A, 1B and 1C are flow cytometric analyses of F1-APC (fluorescent labelled activated protein C) binding to HUVEC (human umbilical vein endothelial cells). Figure 1A is a graph of cell number versus log of fluorescence intensity, demonstrating F1-APC binding to HUVEC. HUVEC (1×10^5) were incubated at room temperature without (dotted line) or with 160 nM of F1-APC (solid line) in the presence of 1.3 mM CaCl_2 . After washing, bound APC was analyzed by flow cytometry. Figure 1B is a graph of fluorescence intensity versus F1-APC concentration (nM) demonstrating the concentration dependence of F1-APC binding to HUVEC. HUVEC were incubated with F1-APC in the absence (open circles) or presence of 1.3 mM CaCl_2 (closed circles) and binding was measured as in A. Mean channel fluorescence intensity is plotted for each F1-APC concentration (between 0 and 800 nM). Figure 1C is a graph of the percent of mean fluorescence versus unlabeled protein concentration ($\mu\text{g/ml}$), demonstrating the effects of unlabeled proteins on F1-APC binding to HUVEC. F1-APC binding to HUVEC was carried out in the presence of the indicated concentrations (between 0 and 100 $\mu\text{g/ml}$) of unlabeled APC, protein C, protein S, factor X and Xa or recombinant Gla-domainless protein C (rGDPC). Figures 2A, 2B, 2C and 2D are graphs of ^{125}I -APC Binding to HUVEC Monolayers. Figure 2A is a graph of the bound APC (cpm $\times 10^{-3}$) versus time (min), showing the time course of ^{125}I -APC binding to HUVEC. HUVEC monolayers (1.2×10^5 cells) were incubated at 4°C with 32 nM (filled squares) or 8

nM (open squares) ^{125}I -APC. At the indicated times, cells were washed and bound radioactivity was measured. Figure 2B is a graph of bound APC (cpm $\times 10^{-3}$) versus unlabeled protein (nM) demonstrating the effects of unlabeled APC and rGDPC on ^{125}I -APC binding to HUVEC. HUVEC were incubated at 4°C for one hour with ^{125}I -labeled APC in the presence of the indicated concentrations (between 0.1 and approximately 1000 nM) of unlabeled APC (open

circles) or rGDPC (closed circles). After washing, bound radioactivity was measured. Figure 2C is a

graph of bound APC (fmol/well) versus free APC (nM) demonstrating the concentration dependence of ^{125}I -APC binding to HUVEC. Monolayers of HUVEC were

incubated with the concentrations of ^{125}I -APC indicated as described above. Specific binding was determined as described below. Figure 2D is a

Scatchard analysis of ^{125}I -APC binding to HUVEC.

Each value was calculated from the data shown in Figure 2C.

Figures 3A and 3B are flow cytometric analyses of F1-APC binding to 293T cells

transfected with a cDNA clone of EPCR. Cells were transfected with a clone EPCR/pEF-BOS or pEF-BOS (negative control) by the calcium/phosphate method.

After 24 h, cells were harvested and F1-APC binding was performed in the absence (dotted lines) or presence of 1.3 mM CaCl_2 (solid lines).

Figures 4a-4c are the predicted protein structure of EPCR based on nucleotide sequence (SEQ ID NO. 1), predicted amino acid sequence (SEQ ID NO. 2) and a hydropathy plot of EPCR. The signal sequence and transmembrane region are indicated with the solid bars.

Figures 5a-5b are a comparison of the amino acid sequence of EPCR to the amino acid sequences of other members of the CD1 family and CCD41. The EPCR sequence is shown in the first line and

compared to murine CCD41 (second line), human CD1d (third line) and murine CD1.2 (fourth line).

Identities with EPCR are indicated by open boxes.

Residues that are conserved between EPCR and all of the human CD1 family members are indicated by a double asterisk. Residues shared with one or more members of the CD1 family are indicated by a single asterisk.

Figure 6 is a comparison of the amino acid sequence of human EPCR (first line) to the amino acid sequence of murine EPCR (second line). Identities are indicated by lines. Similarities are indicated with dots.

Detailed Description of the Invention

I. Cloning and Characterization of EPCR.

Human protein C and activated protein C are shown to bind to endothelium specifically, selectively and saturably ($K_d = 30$ nM, 7000 sites per cell) in a Ca^{2+} dependent fashion. FL-APC

binding to various human cell lines were examined, and found that the binding was HUVEC specific. A human kidney cell line transformed with SV40 large T antigen, 293T cells, expressed very few of these binding sites. A HUVEC cDNA library was

constructed using the powerful mammalian expression vector, pEF-BOS (Mizushima and Nagata, (1990)

Nucleic Acids Res. 18, 5322). Plasmid DNA was prepared from subpools of independent colonies (2,500 colonies per pool), and transfected into

293T cells, using the method of Kaisho et al., (1994) Proc. Natl. Acad. Sci. (USA) 91, 5325.

FL-APC binding was analyzed on a flow cytometer.

One of eight subpools gave a positive signal. This subpool was divided into 20 subpools and

rescreened. After three rounds of enrichment, one positive clone, EPCR-1, was isolated. EPCR-1

carries a 1.3 kb insert. When transfected into 293T cells, this clone was capable of expressing the calcium-dependent binding site for FL-APC on the 293T cell surface.

5 Expression cloning revealed a 1.3 kb cDNA that coded for a type I transmembrane glycoprotein capable of binding protein C. This protein appears to be a member of the CD1/MHC superfamily. Like thrombomodulin, the receptor involved in protein C
10 activation, the endothelial cell protein C receptor (EPCR) function and message are both down regulated by exposure of endothelium to TNF. Identification of EPCR as a member of the CD1/MHC superfamily provides insights into the role of this receptor
15 for protein C in regulating the inflammatory response.

Materials and Methods

Protein Preparation

Human protein C (Esmon et al., (1993)
20 Meths. Enzymol. 222, 359-385), APC (Esmon et al., 1993), recombinant gla domainless protein C (rGDPC) (Rezaie et al., (1992) J. Biol. Chem. 267, 11701-11704), protein S (Taylor et al., 1991), factor X and factor Xa (Le Bonniec et al., (1992)
25 J. Biol. Chem. 267, 6970-6976) were prepared as described in the cited publications.

Selective labeling of the active site of APC with fluorescein was performed by the method of Bock (Bock, P.E. (1988) Biochemistry 27,
30 6633-6639). In brief, N^ε-[(acetylthio)acetyl]-D-Phe-Pro-Arg-CH₂Cl (200 μM) was reacted with 40 μM APC for 1 hour at room temperature. After dialysis, the covalently modified APC was incubated at room temperature for
35 one hour with 200 μM 5-(iodoacetamido) fluorescein (Molecular Probes). Free fluorescein was removed by gel filtration on a PD-10 column (Pharmacia).

With this method, (each molecule of fluoresceinated APC (FI-APC) contains a single dye at the active site and hence all of the fluorescent molecules behave identically.

5 Iodogen (Pierce) was used to radiolabel APC with $\text{Na}[^{125}\text{I}]$ (Amersham) according to the manufacture's protocol in the presence of 5 mM CaCl_2 . Free ^{125}I was removed by gel filtration on a PD-10 column. The specific activity of the ^{125}I -APC was 1×10^4 cpm/ng protein.

Cell Culture

Human umbilical vein endothelial cells (HUVEC) were isolated from fresh umbilical-cords by collagenase treatment and cultured in medium 199 containing 15% fetal bovine serum, 10 $\mu\text{g/ml}$ heparin, and 0.5% endothelial cell growth supplement prepared from bovine brain extract (Maciag et al., (1979) Proc. Natl. Acad. Sci. (USA) 76, 5674-5678). HOS (ATCC CRL 1543), Hep-2 (ATCC CCL 23) and 293 cells (ATCC CRL 1573) transformed with SV40 large T antigen (293T, a gift from Dr. Kenji Oritani) were maintained in Earl's MEM supplemented with 10% fetal bovine serum. The human lymphocyte cell lines, Jurkat, MOLT3 (ATCC CRL 1552), Jijoye (ATCC CCL 87), Raji (ATCC CCL 86), U-937 (ATCC CRL 1593), HL-60 (ATCC CCL 240), and HEL (ATCC TIB 180), were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum.

Flow Cytometric Analysis of FI-APC Binding to Cells

Adherent cells were harvested by incubation at 37°C for 5 min in phosphate buffered saline (PBS) containing 0.02% EDTA. Cells were washed twice with EDTA/PBS and then once with Hank's balanced salt solution (HBSS). They were resuspended in HBSS containing 1% bovine serum albumin (BSA) and 0.02% sodium azide (binding

10

¹²⁵I-APC binding to HUVEC

15

Construction of HUVEC CDNA Library

30

Nagata, 1990; this vector was a kind gift from Dr. S. Nagata). The construct was transfected into *E. coli* DH10B by electroporation (Bio-Rad Gene Pulser™). The library- consisted of 8×10^6 independent colonies with an average size of 2.0 kb.

Expression Cloning and Sequence Analysis

Approximately 2×10^6 independent colonies were divided into eight subpools (each containing 2,500 independent colonies) and plasmid DNA was prepared from each subpool. Sub-confluent 293T cells in 24-well microplates were transfected with 1 μ g of the DNA by the calcium/phosphate method (Graham and Van Der Eb, (1973) *Virology* 52, 456-467). After 20 hours, the medium was changed, and culture was continued for another 24 hours. The subpools were screened for F1-APC binding by FACS analysis as described above. The positive library pool was then divided into 20 new pools and rescreened. After three rounds of screening, 96 individual clones were tested and one positive clone was identified.

The insert (1.3 kb) was subcloned into pBluescript™ (Stratagene), and the nucleotide sequence was determined using a Sequenase™ version 2.0 DNA Sequencing kit (USB). Nucleotide and protein database search employed the BLAST™ (NCBI) and FASTA™ programs (GCG) with GenBank, EMBL, and SwissProt databases.

Northern Blot Analysis

Total RNAs (15 μ g) from various cells were isolated, electrophoresed through formaldehyde agarose gels and transferred to a nylon membrane (Hybond-N™, Amersham). The 483 bp *Xba* I fragment from the 5' end of the EPCR cDNA was labeled by random priming according to the manufacturer's

instructions (Multiprime™ DNA labeling system, Amersham) and used for hybridization.

Protein C and APC Binding to HUVEC

Endothelial cells in suspension bound FL-APC, as monitored by flow cytometry, and demonstrated in Figure 1A. Binding was saturable and Ca^{2+} dependent, as shown by Figure 1B. Optimal binding required at least 1 mM Ca^{2+} . FL-APC was displaced from the cell surface by APC and protein C equivalently, as shown by Figure 1C. The homologous Gla-domain containing proteins, protein S, factor X, and its active form, factor Xa, failed to displace bound F1-APC, suggesting that there is a specific binding site for APC on the endothelial cell surface. Protein C binding was dependent on the Gla domain, since recombinant gla-domainless protein C (rGDPC) failed to displace F1-APC.

Detailed binding studies were also performed with ^{125}I -labeled APC and monolayers of HUVEC, as shown by Figures 2A, 2B, 2C and 2D. The binding analysis indicated 7,000 sites per cell and a $K_d=30$ nM. This affinity is similar to that estimated from Figure 1.

Endothelial cell surface thrombomodulin can interact with protein C and APC. The K_d (greater than 1 μM) (Hogg et al., (1992) J. Biol. Chem. 267, 703-706; Olsen et al., (1992) Biochemistry 31, 746-754), however, is much higher than that of the binding site described above with respect to the new receptor. Furthermore, polyclonal and monoclonal antibodies against thrombomodulin that inhibit protein C activation did not inhibit the binding. Protein S also can interact with protein C and APC (Dahlbäck et al., (1992) Biochemistry 31, 12769-12777), but F1-APC binding to HUVEC was not influenced by protein S addition. Furthermore, polyclonal and monoclonal

antibodies to protein S did not inhibit the binding. These results indicate the binding site for protein C and APC on endothelium is distinct from these known molecules.

5

Nucleotide and Predicted Protein Structure Analysis of EPCR

The insert was subcloned into pBluescript, and the nucleotide sequence was determined, as shown in Sequence ID No. 1. The
10 cDNA shown in Sequence ID No. 1 consists of 1302 bp, including a translation initiation ATG codon (AGGATGT, (Kozak, (1986) Cell 44, 283-292) at the 5'-end at nucleotides 25-27 of Sequence ID No. 1. A potential polyadenylation signal sequence,
15 AATAAA, (Proudfoot and Brownlee, (1976) Nature 263, 211-214) begins at nucleotide 1267 of Sequence ID No. 1, just 18 bp upstream of the poly(A) sequence.

The cDNA is predicted to code for a protein of 238 amino acids (Sequence ID No. 2),
20 which includes a 15 amino acid signal sequence (von Heijne, (1986) Nucleic Acids Res. 14, 4683-4690) at the N-terminal. Therefore, the mature protein is predicted to contain 223 amino acids. Sequence ID No. 2 is the predicted amino acid sequence of EPCR.
25 Amino acids 1-15 of Sequence ID No. 2

(MLTTLLPILLLSGWA) are the putative signal sequence determined by the method of von Heijne (von Heijne, 1986). Amino acids 211-236 of Sequence ID No. 2 (LVLGVLVGGFIIAGVAVGIFLCTGGR) are the putative
30 transmembrane domain. Potential N-glycosylation sites are present at amino acids 47-49, 64-66, 136-138, and 172-174 of Sequence ID No. 2.

Extracellular cysteine residues are present at amino acids 17, 114, 118, and 186 of Sequence ID
35 No. 2. A potential transmembrane region (Engelman et al., (1986) Annu. Rev. Biophys. Chem. 15, 321-53) consisting of 23 amino acids was identified at

the C-terminal end (beginning at amino acid 216 of Sequence ID No. 2).

The protein is predicted to be a type 1 transmembrane protein. The extracellular domain contains four potential N-glycosylation sites and four Cys residues. The cytoplasmic region contains only three amino acids and terminates with a Cys, which could be acylated to something or involved in heterodimer formation with another peptide.

Although described with reference to cloning and expression of the protein encoding sequence, larger amounts of protein can be obtained by expression in suitable recombinant host systems, such as mammalian, yeast, bacteria, or insect cells. Isolation can be facilitated by making antibodies to the recombinant protein which are then immobilized on substrates for use in purification of additional receptors, as described below.

As used herein, the nucleotide sequences encoding the receptor include the sequence shown in Sequence ID No. 1, and sequences having conservative substitutions, additions or deletions thereof which hybridize to Sequence ID No. 1 under stringent conditions. As used herein, the amino acids sequences constituting the receptor include the sequence shown in Sequence ID No. 2, and sequences having conservative substitutions, additions or deletions thereof which form a receptor having functionally equivalent biological activity. It is well known to those skilled in the art what constitutes conservative substitutions, additions or deletions, and which could be readily ascertained as encoding, or forming, a functionally equivalent receptor molecule using the functional assays described herein.

The hydropathic plot shown in Figures 4a-4c was performed according to the method of Goldman et al (Engelman et al., 1994) (solid line) and that of Kyte and Doolittle (1982) (J. Mol. Biol. 157, 332-50; 105-132) (dotted line).

DNA and protein database searches revealed that the sequence is related to the centrosome-associated, cell cycle dependent murin protein, CCD41, also referred as centrocyclin (Rothbarth et al., (1993) J. Cell Sci. 104, 19-30), as shown by Figures 5a-5b. The similarity in the published sequence of murine CCD41 with human EPCR led to the cloning and sequencing of the murine EPCR. The sequence of murine EPCR is shown in Figure 6. It is distinct from the published sequence of CCD41.

The EPCR amino acid sequence was also related to, but quite distinct from, the CD1/MHC superfamily and the murine CD1.2, as also shown by Figures 5a-5b. Based on the homology to CD1/MHC, it is likely that EPCR contains two domains consisting of residues 17-114 and 118-188. Of the CD1 family members, CD1d is the most similar to EPCR. In the mouse, CCD41 is associated exclusively with the centrosome during G₁ but becomes detectable elsewhere during the cell cycle, reaching a maximum during G₂, except during the G₂/M phase (Rothbarth et al., 1993). EPCR expression appears restricted to endothelium, which would not be expected for a cell cycle associated protein.

The identification of the protein C receptor on endothelium suggests that the endothelial cell binds protein C/APC through three distinct mechanisms. In addition to EPCR, protein S can bind APC/protein C on negatively charged membrane surfaces that include the endothelium (Stern et al., (1986) J. Biol. Chem. 261, 713-718),

but this is not cell-type specific (Dahlbäck et al., 1992). Thrombomodulin in complex with thrombin can bind protein C and APC (Hogg et al., 1992). On endothelium, the protein S binding sites (Nawroth and Stern, (1986) J. Exp. Med. 163, 740-745), thrombomodulin (Esmon, 1989) and EPCR are all down regulated by cytokines, indicating that inflammation can impair protein C pathway function at multiple levels.

10 The homology to the CD1/MHC family of proteins is especially interesting since it provides indications as to the function of EPCR. The CD1/MHC family has three extracellular domains termed $\alpha 1$, $\alpha 2$ and $\alpha 3$. The extracellular domain of EPCR 15 contains four Cys residues that appear to correspond to two distinct domains. EPCR lacks the third domain of the CD1/MHC family, but the two domains have significant homology to the $\alpha 1$ and $\alpha 2$ domains of the CD1 protein family and the $\alpha 2$ domain 20 of the MHC class II protein, suggesting that these proteins evolved from a common ancestor. The first domain of EPCR, residues 17-114, contains two potential N glycosylation sites and is rich in β strand structure, suggesting that it may form a β 25 sheet. Despite the β strand structure, consensus sequences (Williams and Barclay, (1988) Ann. Rev. Immunol. 6, 381-405) for the immunoglobulin superfamily of receptors are absent. The second domain of EPCR, residues 118-188, contains two 30 additional N glycosylation sites and, like the CD1/MHC family, this domain is predicted to have limited β structure.

II. Modulation of Inflammation using EPCR.

In vitro studies have suggested 35 anti-inflammatory activities for APC. For instance, an unusual carbohydrate sequence on protein C can inhibit inflammatory cell adhesion to

selectins (Grinnell et al., (1994) Glycobiology, 4, 221-226). Modest inhibitory effects of APC have been reported on TNF production (Grey et al., (1993) Transplant. Proc. 25, 2913-2914). EPCR could contribute to these anti-inflammatory mechanisms. Since the homologous protein family, CD1, can be linked to CD8 (Ledbetter et al., (1985) J. Immunol. 134, 4250-4254), it is also possible that the protein C receptor is linked to another protein and signal through this second protein. One of the CD1 family members, CD1d, has been reported to promote T cell responses, possibly involving binding to CD8 (Panja et al., (1993) J. Exp. Med. 178, 1115-1119). CD1b has recently been reported to serve as an antigen-presenting molecule (Porcelli et al., (1992) Nature 360, 593-597). The ability to bind protein C/APC could then be linked either directly or indirectly to signalling via direct interaction with cells of the immune system. Since the MHC class of proteins is involved in presentation of proteins to cell receptors, the concept of presentation of protein C/APC to inflammatory cells as a means of elaborating anti-inflammatory activity may also be involved. This includes modulation of enzyme specificity such as occurs with thrombin-thrombomodulin interaction (Esmon, 1989). In this case, the EPCR-APC complex might cleave biologically active peptides from unknown substrates.

EPCR mRNA Levels and APC Binding

To determine the cellular specificity of EPCR expression, the intensity of FL-APC binding to HUVEC was compared to several human cell lines. FL-APC bound strongly only to HUVEC, and not to any of the T, B, or monocytic cell lines tested. Cells were incubated at room temperature without or with 160 nM FL-APC in the presence of 1.3 mM CaCl_2 .

Binding was analyzed by flow cytometry. Slight binding was demonstrated with the osteosarcoma line, HOS and the epidermoid carcinoma cell line, HEP-2.

5 Total RNA was extracted from these cells and hybridized with the EPCR cDNA probe for Northern Blot Analysis. EPCR mRNA was detected by Northern blot analysis for HUVEC, Jurkat, HEP-2, Raji, HOS, and U937. Among the cells lines tested, 10 EPCR mRNA was detected at high levels only in HUVEC. The calculated mRNA size of 1.3 kb was identical to the size of the isolated cDNA. After prolonged exposure, a weak signal was also detected with the osteosarcoma cell line HOS and monocyte 15 cell line U937. Thus, both APC binding and EPCR mRNA expression are very specific for endothelium.

Effects of TNF on APC Binding and EPCR MRNA Levels

Several other members of the protein C 20 anticoagulant pathway are subject to regulation by inflammatory cytokines (Esmon, 1989). For instance, endothelial cell surface thrombomodulin expression and message are known to be reduced by exposure of the cells to TNF (Conway and Rosenberg, 25 1988; Lentz et al., 1991). To determine if a similar process occurs with EPCR, HUVEC were treated with TNF and APC binding and expression of EPCR mRNA were examined. APC binding to HUVEC decreased in a time dependent fashion. EPCR 30 activity decreased more rapidly than thrombomodulin antigen. HUVEC were cultured for 0, 6, 24 and 48 hr, in the presence of TNF- α (10 ng/ml). Cells were harvested and residual F1-APC binding or thrombomodulin (TM) expression was analyzed by flow 35 cytometry. Cell surface TM was stained with an anti-TM murine monoclonal antibody and FITC--conjugated anti-mouse IgG. The negative control is without added fluorescent ligand.

HUVEC were treated with 10 ng/ml of TNF- α for 0, 0.5, 1, 2, 3, 6, 10 and 24 hr, and message was extracted and detected as described above. The results demonstrated that the concentration of EPCR mRNA was also reduced by TNF treatment. Message levels and APC binding activity decreased in parallel. Therefore, the TNF mediated down-regulation of APC binding to endothelium probably occurs at the level of mRNA expression.

Enhancement of inflammatory responses by blocking binding of endogenous molecules to EPCR can be achieved by administration of compounds binding to the receptor to a subject in need of inhibition. The degree of binding is routinely determined using assays such as those described above. Compounds which are effective include antibodies to the protein, fragments of antibodies retaining the binding regions, and peptide fragments of APC which include the Gla region.

Inhibition of the inflammatory response could be obtained by infusing soluble EPCR. Alternatively, localizing EPCR to surfaces in contact with blood would render the surfaces anticoagulant by virtue of the ability of EPCR to bind and concentrate the anticoagulant APC at the surface. Alternatively, the function of EPCR could be enhanced by overexpressing the EPCR in endothelium used to coat vascular grafts in patients with vascular disease or on stents in cardiac patients.

The DNA sequence can also be used for screening for other homologous or structurally similar receptor proteins using hybridization probes.

These methods and reagents and pharmaceuticals are more readily understood by reference to the following.

Screening of patient samples for
expression of receptor proteins.

Patients with thrombosis or hyperinflammatory conditions could be screened for defects in the EPCR gene. Sequence ID No. 1, and consecutive portions thereof of at least about seven nucleotides, more preferably fourteen to seventeen nucleotides, most preferably about twenty nucleotides, are useful in this screening using hybridization assays of patient samples, including blood and tissues. Screening can also be accomplished using antibodies, typically labelled with a fluorescent, radiolabelled, or enzymatic label, or by isolation of target cells and screening for binding activity, as described in the examples above. Typically, one would screen for expression on either a qualitative or quantitative basis, and for expression of functional receptor. Labelling can be with ^{32}P , ^{35}S , fluorescein, biotin, or other labels routinely used with methods known to those skilled in the art for labelling of proteins and/or nucleic acid sequences.

Hybridization Probes

Reaction conditions for hybridization of an oligonucleotide probe or primer to a nucleic acid sequence vary from oligonucleotide to oligonucleotide, depending on factors such as oligonucleotide length, the number of G and C nucleotides, and the composition of the buffer utilized in the hybridization reaction. Moderately stringent hybridization conditions are generally understood by those skilled in the art as conditions approximately 25°C below the melting temperature of a perfectly base-paired double-stranded DNA. Higher specificity is generally achieved by employing incubation conditions having higher temperatures, in other words, more stringent conditions. In general, the longer the sequence

or higher the G and C content, the higher the temperature and/or salt concentration required.

Chapter 11 of the well-known laboratory manual of Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, second edition, Cold Spring Harbor Laboratory Press, New York (1990), (which is incorporated by reference herein), describes hybridization conditions for oligonucleotide probes and primers in great detail, including a description of the factors involved and the level of stringency necessary to guarantee hybridization with specificity.

The preferred size of a hybridization probe is from 10 nucleotides to 100,000 nucleotides in length. Below 10 nucleotides, hybridized systems are not stable and will begin to denature above 20°C. Above 100,000 nucleotides, one finds that hybridization (renaturation) becomes a much slower and incomplete process, as described in greater detail in the text MOLECULAR GENETICS, Stent, G.S. and R. Calender, pp. 213-219 (1971). Ideally, the probe should be from 20 to 10,000 nucleotides. Smaller nucleotide sequences (20-100) lend themselves to production by automated organic synthetic techniques. Sequences from 100-10,000 nucleotides can be obtained from appropriate restriction endonuclease treatments. The labeling of the smaller probes with the relatively bulky chemiluminescent moieties may in some cases interfere with the hybridization process.

Generation of Antibodies for Diagnostic or Therapeutic Use

Antibodies to the receptor proteins can also be generated which are useful in detection, characterization or isolation of receptor proteins, as well as for modifying receptor protein activity, in most cases, through inhibition of binding.

Antibodies are generated by standard techniques,

using human or animal receptor proteins. Since the proteins exhibit high evolutionary conservation, it may be advantageous to generate antibodies to a protein of a different species of origin than the species in which the antibodies are to be tested or utilized, looking for those antibodies which are immunoreactive with the most evolutionarily conserved regions. Antibodies are typically generated by immunization of an animal using an adjuvant such as Freund's adjuvant in combination with an immunogenic amount of the protein administered over a period of weeks in two to three week intervals, then isolated from the serum, or used to make hybridomas which express the antibodies in culture. Because the methods for immunizing animals yield antibody which is not of human origin, the antibodies could elicit an adverse effect if administered to humans. Methods for "humanizing" antibodies, or generating less immunogenic fragments of non-human antibodies, are well known. A humanized antibody is one in which only the antigen-recognized sites, or complementarily-determining hypervariable regions (CDRs) are of non-human origin, whereas all framework regions (FR) of variable domains are products of human genes. These "humanized" antibodies present a lesser xenographic rejection stimulus when introduced to a human recipient.

To accomplish humanization of a selected mouse monoclonal antibody, the CDR grafting method described by Daugherty, et al., (1991) Nucl. Acids Res., 19:2471-2476, incorporated herein by reference, may be used. Briefly, the variable region DNA of a selected animal recombinant anti-idiotypic ScFv is sequenced by the method of Clackson, T., et al., (1991) Nature, 352:624-688, incorporated herein by reference. Using this

sequence, animal CDRs are distinguished from animal framework regions (FR) based on locations of the CDRs in known sequences of animal variable genes. Kabat, H.A., et al., Sequences of Proteins of Immunological Interest, 4th Ed. (U.S. Dept. Health and Human Services, Bethesda, MD, 1987). Once the animal CDRs and FR are identified, the CDRs are grafted onto human heavy chain variable region framework by the use of synthetic oligonucleotides and polymerase chain reaction (PCR) recombination. Codons for the animal heavy chain CDRs, as well as the available human heavy chain variable region framework, are built in four (each 100 bases long) oligonucleotides. Using PCR, a grafted DNA sequence of 400 bases is formed that encodes for the recombinant animal CDR/human heavy chain FR protection.

The immunogenic stimulus presented by the monoclonal antibodies so produced may be further decreased by the use of Pharmacia's (Pharmacia LKB Biotechnology, Sweden) "Recombinant Phage Antibody System" (RPAS), which generates a single-chain Fv fragment (ScFv) which incorporates the complete antigen-binding domain of the antibody. In the RPAS, antibody variable heavy and light chain genes are separately amplified from the hybridoma mRNA and cloned into an expression vector. The heavy and light chain domains are co-expressed on the same polypeptide chain after joining with a short linker DNA which codes for a flexible peptide. This assembly generates a single-chain Fv fragment (ScFv) which incorporates the complete antigen-binding domain of the antibody. Compared to the intact monoclonal antibody, the recombinant ScFv includes a considerably lower number of epitopes, and thereby presents a much weaker immunogenic stimulus when injected into humans.

The antibodies can be formulated in standard pharmaceutical carriers for administration to patients in need thereof. These include saline, phosphate buffered saline, and other aqueous carriers; and liposomes, polymeric microspheres and other controlled release delivery devices, as are well known in the art. The antibodies can also be administered with adjuvant, such as muramyl dipeptide or other materials approved for use in humans (Freund's adjuvant can be used for administration of antibody to animals).

Screening for drugs modifying or altering the extent of receptor function or expression

The receptor proteins are useful as targets for compounds which turn on, or off, or otherwise regulate binding to these receptors. The assays described above clearly provide routine methodology by which a compound can be tested for an inhibitory effect on binding of PC or APC. The *in vitro* studies of compounds which appear to inhibit binding selectively to the receptors are then confirmed by animal testing. Since the molecules are so highly evolutionarily conserved, it is possible to conduct studies in laboratory animals such as mice to predict the effects in humans.

In cases where inflammatory mediators or vascular disease down regulate EPCR, it would be advantageous to increase its concentration *in vivo* on endothelium. The binding assays described here and the gene sequence allow assays for increased EPCR expression. Similar approaches have been taken with thrombomodulin and investigators have shown that cyclic AMP (Maruyama, I. et al. (1991) Thrombosis Research 61, 301-310) and interleukin 4 (Kapiotis, S. et al., (1991) Blood 78, 410-415) can elevate thrombomodulin expression. The ability to

screen such drugs or drugs that block TNF down regulation of EPCR provide an approach to elevating EPCR expression *in vivo* and thus enhancing anticoagulant and anti-inflammatory activity.

5 Studies based on inhibition of binding are predictive for indirect effects of alteration of receptor binding. For example, inhibition of binding of APC or increased expression of TNF is predictive of inhibition of EPCR function.

10 Assays for testing compounds for useful activity can be based solely on interaction with the receptor protein, preferably expressed on the surface of transfected cells such as those described above. Proteins in solution or
15 immobilized on inert substrates can also be utilized. These can be used to detect inhibition or enhancement in binding of PC or APC

 Alternatively, the assays can be based on interaction with the gene sequence encoding the
20 receptor protein, preferably the regulatory sequences directing expression of the receptor protein. For example, antisense which binds to the regulatory sequences, and/or to the protein encoding sequences can be synthesized using
25 standard oligonucleotide synthetic chemistry. The antisense can be stabilized for pharmaceutical use using standard methodology (encapsulation in a liposome or microsphere; introduction of modified nucleotides that are resistant to degradation or
30 groups which increase resistance to endonucleases, such as phosphorothiodates and methylation), then screened initially for alteration of receptor activity in transfected or naturally occurring cells which express the receptor, then *in vivo* in
35 laboratory animals. Typically, the antisense would inhibit expression. However, sequences which block

those sequences which "turn off" synthesis can also be targeted.

The receptor protein for study can be isolated from either naturally occurring cells or cells which have been genetically engineered to express the receptor, as described in the examples above. In the preferred embodiment, the cells would have been engineered using the intact gene.

Random generation of receptor or receptor encoding sequence binding molecules.

Molecules with a given function, catalytic or ligand-binding, can be selected for from a complex mixture of random molecules in what has been referred to as "in vitro genetics" (Szostak, (1992) TIBS 19:89). One synthesizes a large pool of molecules bearing random and defined sequences and subjects that complex mixture, for example, approximately 10^{15} individual sequences in 100 μ g of a 100 nucleotide RNA, to some selection and enrichment process. For example, by repeated cycles of affinity chromatography and PCR amplification of the molecules bound to the ligand on the column, Ellington and Szostak (1990) estimated that 1 in 10^{10} RNA molecules folded in such a way as to bind a given ligand. DNA molecules with such ligand-binding behavior have been isolated (Ellington and Szostak, 1992; Bock et al, 1992).

30

Computer assisted drug design

Computer modeling technology allows visualization of the three-dimensional atomic structure of a selected molecule and the rational design of new compounds that will interact with the molecule. The three-dimensional construct typically depends on data from x-ray crystallographic analyses or NMR imaging of the selected molecule. The molecular dynamics require

35

force field data. The computer graphics systems enable prediction of how a new compound will link to the target molecule and allow experimental manipulation of the structures of the compound and target molecule to perfect binding specificity.

Prediction of what the molecule-compound interaction will be when small changes are made in one or both requires molecular mechanics software and computationally intensive computers, usually coupled with user-friendly, menu-driven interfaces between the molecular design program and the user.

Examples of molecular modelling systems are the CHARMM and QUANTA programs, Polygen Corporation, Waltham, MA. CHARMM performs the energy minimization and molecular dynamics functions. QUANTA performs the construction, graphic modelling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with each other.

A number of articles review computer modeling of drugs interactive with specific proteins, such as Rotivinen, et al., (1988) Acta Pharmaceutica Fennica 97, 159-166; Ripka, New Scientist 54-57 (June 16, 1988); McKinaly and Rossmann, (1989) Annu. Rev. Pharmacol. Toxicol. 29, 111-122; Perry and Davies, QSAR: Quantitative Structure-Activity Relationships in Drug Design pp. 189-193 (Alan R. Liss, Inc. 1989); Lewis and Dean, (1989) Proc. R. Soc. Lond. 236, 125-140 and 141-162; and, with respect to a model receptor for nucleic acid components, Askew, et al., (1989) J. Am. Chem. Soc. 111, 1082-1090. Other computer programs that screen and graphically depict chemicals are available from companies such as BioDesign, Inc., Pasadena, CA., Allelix, Inc., Mississauga, Ontario, Canada, and Hypercube, Inc.,

Cambridge, Ontario. Although these are primarily designed for application to drugs specific to particular proteins, they can be adapted to design of drugs specific to regions of DNA or RNA, once that region is identified.

Although described above with reference to design and generation of compounds which could alter binding, one could also screen libraries of known compounds, including natural products or synthetic chemicals, and biologically active materials, including proteins, for compounds which are inhibitors or activators.

Generation of nucleic acid regulators

Nucleic acid molecules containing the 5' regulatory sequences of the receptor genes can be used to regulate or inhibit gene expression in vivo. Vectors, including both plasmid and eukaryotic viral vectors, may be used to express a particular recombinant 5' flanking region-gene construct in cells depending on the preference and judgment of the skilled practitioner (see, e.g., Sambrook et al., Chapter 16). Furthermore, a number of viral and nonviral vectors are being developed that enable the introduction of nucleic acid sequences in vivo (see, e.g., Mulligan, (1993) Science, 260, 926-932; United States Patent No. 4,980,286; United States Patent No. 4,868,116; incorporated herein by reference). Recently, a delivery system was developed in which nucleic acid is encapsulated in cationic liposomes which can be injected intravenously into a mammal. This system has been used to introduce DNA into the cells of multiple tissues of adult mice, including endothelium and bone marrow (see, e.g., Zhu et al., (1993) Science 261, 209-211; incorporated herein by reference).

The 5' flanking sequences of the receptor gene can also be used to inhibit the expression of the receptor. For example, an antisense RNA of all or a portion of the 5' flanking region of the receptor gene can be used to inhibit expression of the receptor in vivo. Expression vectors (e.g., retroviral expression vectors) are already available in the art which can be used to generate an antisense RNA of a selected DNA sequence which is expressed in a cell (see, e.g., U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286). Accordingly, DNA containing all or a portion of the sequence of the 5' flanking region of the receptor gene can be inserted into an appropriate expression vector so that upon passage into the cell, the transcription of the inserted DNA yields an antisense RNA that is complementary to the mRNA transcript of the receptor protein gene normally found in the cell. This antisense RNA transcript of the inserted DNA can then base-pair with the normal mRNA transcript found in the cell and thereby prevent the mRNA from being translated. It is of course necessary to select sequences of the 5' flanking region that are downstream from the transcriptional start sites for the receptor protein gene to ensure that the antisense RNA contains complementary sequences present on the mRNA. Antisense RNA can be generated in vitro also, and then inserted into cells. Oligonucleotides can be synthesized on an automated synthesizer (e.g., Model 8700 automated synthesizer of Milligen-Bioscience, Burlington, MA or ABI Model 380B). In addition, antisense deoxyoligonucleotides have been shown to be effective in inhibiting gene transcription and viral replication (see e.g., Zamecnik et al.,

(1978) Proc. Natl. Acad. Sci. USA 75, 280-284; Zamecnik et al., (1986) Proc. Natl. Acad. Sci., 83, 4143-4146; Wickstrom et al., (1988) Proc. Natl. Acad. Sci. USA 85, 1028-1032; Crooke, (1993) FASEB J. 7, 533-539. Furthermore, recent work has shown that improved inhibition of expression of a gene by antisense oligonucleotides is possible if the antisense oligonucleotides contain modified nucleotides (see, e.g., Offensperger et al., (1993) EMBO J. 12, 1257-1262 (in vivo inhibition of duck hepatitis B viral replication and gene expression by antisense phosphorothioate oligodeoxynucleotides); PCT WO 93/01286 Rosenberg et al., (synthesis of sulfurthioate oligonucleotides); Agrawal et al., (1988) Proc. Natl. Acad. Sci. USA 85, 7079-7083 (synthesis of antisense oligonucleoside phosphoramidates and phosphorothioates to inhibit replication of human immunodeficiency virus-1); Sarin et al., (1989) Proc. Natl. Acad. Sci. USA 85, 7448-7794 (synthesis of antisense methylphosphonate oligonucleotides); Shaw et al., (1991) Nucleic Acids Res. 19, 747-750 (synthesis of 3' exonuclease-resistant oligonucleotides containing 3' terminal phosphoroamidate modifications); incorporated herein by reference). The sequences of the 5' flanking region of receptor protein gene can also be used in triple helix (triplex) gene therapy. Oligonucleotides complementary to gene promoter sequences on one of the strands of the DNA have been shown to bind promoter and regulatory sequences to form local triple nucleic acid helices which block transcription of the gene (see, e.g., Maher et al., (1989) Science 245, 725-730; Orson et al., (1991) Nucl. Acids Res. 19, 3435-3441; Postal et al., (1991) Proc. Natl. Acad. Sci. USA 88, 8227-8231;

Cooney et al., (1988) Science 241, 456-459; Young et al., (1991) Proc. Natl. Acad. Sci. USA 88, 10023-10026; Duval-Valentin et al., (1992) Proc. Natl. Acad. Sci. USA 89, 504-508; Blume et al., (1992) Nucl. Acids Res. 20, 1777-1784; Grigoriev et al., (1992) J. Biol. Chem. 267, 3389-3395.

Recently, both theoretical calculations and empirical findings have been reported which provide guidance for the design of oligonucleotides for use in oligonucleotide-directed triple helix formation to inhibit gene expression. For example, oligonucleotides should generally be greater than 14 nucleotides in length to ensure target sequence specificity (see, e.g., Maher et al., (1989); Grigoriev et al., (1992)). Also, many cells avidly take up oligonucleotides that are less than 50 nucleotides in length (see e.g., Orson et al., (1991); Holt et al., (1988) Mol. Cell. Biol. 8, 963-973; Wickstrom et al., (1988) Proc. Natl. Acad. Sci. USA 85, 1028-1032). To reduce susceptibility to intracellular degradation, for example by 3' exonucleases, a free amine can be introduced to a 3' terminal hydroxyl group of oligonucleotides without loss of sequence binding specificity (Orson et al., 1991). Furthermore, more stable triplexes are formed if any cytosines that may be present in the oligonucleotide are methylated, and also if an intercalating agent, such as an acridine derivative, is covalently attached to a 5' terminal phosphate (e.g., via a pentamethylene bridge); again without loss of sequence specificity (Maher et al., (1989); Grigoriev et al., (1992)).

Methods to produce or synthesize oligonucleotides are well known in the art. Such methods can range from standard enzymatic digestion followed by nucleotide fragment isolation (see e.g., Sambrook et al., Chapters 5, 6) to purely

synthetic methods, for example, by the cyanoethyl phosphoramidite method using a Milligen or Beckman System 1Plus DNA synthesizer (see also, Ikuta et al., (1984) Ann. Rev. Biochem. 53, 323-356 (phosphotriester and phosphite-triester methods); Narang et al., (1980) Methods Enzymol., 65, 610-620 (phosphotriester method). Accordingly, DNA sequences of the 5' flanking region of the receptor protein gene described herein can be used to design and construct oligonucleotides including a DNA sequence consisting essentially of at least 15 consecutive nucleotides, with or without base modifications or intercalating agent derivatives, for use in forming triple helices specifically within the 5' flanking region of a receptor protein gene in order to inhibit expression of the gene.

In some cases it may be advantageous to insert enhancers or multiple copies of the regulatory sequences into an expression system to facilitate screening of methods and reagents for manipulation of expression.

Preparation of Receptor Protein Fragments

Compounds which are effective for blocking binding of the receptor can also consist of fragments of the receptor proteins, expressed recombinantly and cleaved by enzymatic digest or expressed from a sequence encoding a peptide of less than the full-length receptor protein. These will typically be soluble proteins, i.e., not including the transmembrane and cytoplasmic regions, although smaller portions determined in the assays described above to inhibit or compete for binding to the receptor proteins can also be utilized. It is a routine matter to make appropriate receptor protein fragments, test for binding, and then utilize. The preferred fragments are of human origin, in order to minimize potential

immunological response. The peptides can be as short as five to eight amino acids in length and are easily prepared by standard techniques. They can also be modified to increase in vivo half-life, by chemical modification of the amino acids or by attachment to a carrier molecule or inert substrate. Based on studies with other peptide fragments blocking receptor binding, the IC_{50} , the dose of peptide required to inhibit binding by 50%, ranges from about $1 \mu M$ to greater than 10 mM, depending on the peptide size and folding. These ranges are well within the effective concentrations for the in vivo administration of peptides, based on comparison with the RGD-containing peptides, described, for example, in U.S. Patent No. 4,792,525 to Ruoslahti, et al., used in vivo to alter cell attachment and phagocytosis. The peptides can also be conjugated to a carrier protein such as keyhole limpet hemocyanin by its N-terminal cysteine by standard procedures such as the commercial Inject kit from Pierce Chemicals or expressed as a fusion protein, which may have increased efficacy.

As noted above, the peptides can be prepared by proteolytic cleavage of the receptor proteins, or, preferably, by synthetic means. These methods are known to those skilled in the art. An example is the solid phase synthesis described by J. Merrifield, (1964) J. Am. Chem. Soc. 85, 2149, used in U.S. Patent No. 4,792,525, and described in U.S. Patent No. 4,244,946, wherein a protected alpha-amino acid is coupled to a suitable resin, to initiate synthesis of a peptide starting from the C-terminus of the peptide. Other methods of synthesis are described in U.S. Patent No. 4,305,872 and 4,316,891. These methods can be used to synthesize peptides having identical

sequence to the receptor proteins described herein, or substitutions or additions of amino acids, which can be screened for activity as described above.

The peptide can also be administered as a pharmaceutically acceptable acid- or base- addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

Peptides containing cyclopropyl amino acids, or amino acids derivatized in a similar fashion, can also be used. These peptides retain their original activity but have increased half-lives in vivo. Methods known for modifying amino acids, and their use, are known to those skilled in the art, for example, as described in U.S. Patent No. 4,629,784 to Stammer. The peptides are generally active when administered parenterally in amounts above about 1 $\mu\text{g/kg}$ of body weight. Based on extrapolation from other proteins, for treatment of most inflammatory disorders, the dosage range will be between 0.1 to 70 mg/kg of body weight. This dosage will be dependent, in part, on whether one or more peptides are administered.

Pharmaceutical Compositions

Compounds which alter receptor protein binding are preferably administered in a pharmaceutically acceptable vehicle. Suitable

pharmaceutical vehicles are known to those skilled in the art. For parenteral administration, the compound will usually be dissolved or suspended in sterile water or saline. For enteral

5 administration, the compound will be incorporated into an inert carrier in tablet, liquid, or capsular form. Suitable carriers may be starches or sugars and include lubricants, flavorings, binders, and other materials of the same nature.

10 The compounds can also be administered locally by topical application of a solution, cream, gel, or polymeric material (for example, a Pluronic™, BASF).

Alternatively, the compound may be

15 administered in liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a patient are known to those skilled in the art. U.S. Patent No. 4,789,734 describe methods for encapsulating

20 biological materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed or sonicated, as necessary. A

25 review of known methods is by G. Gregoriadis, Chapter 14. "Liposomes", Drug Carriers in Biology and Medicine pp. 287-341 (Academic Press, 1979). Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be

30 tailored for passage through the gastrointestinal tract directly into the bloodstream.

Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time,

35 ranging from days to months. See, for example, U.S. Patent Nos. 4,906,474, 4,925,673, and 3,625,214.

Disorders to be treated

As described herein, a variety of compounds can be used to inhibit or enhance expression of the EPCR. The nature of the disorder will determine if the expression should be enhanced or inhibited. For example, based on the studies involving the use of an anti-protein C antibody in combination with cytokine, it should be possible to treat solid tumors by enhancing an inflammatory response involving blocking of protein C or activated protein C binding to an endothelial cell protein C/activated protein C receptor by administering to a patient in need of treatment thereof an amount of a compound blocking binding of protein C or activated protein C to the receptor. Similarly, it should be possible to treat disorders such as gram negative sepsis, stroke, thrombosis, septic shock, adult respiratory distress syndrome, and pulmonary emboli using a method for inhibiting an inflammatory response involving administration of EPCR or EPCR fragments or substances that upregulate EPCR expression to a patient in need of treatment thereof.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Oklahoma Medical Research Foundation
- (ii) TITLE OF INVENTION: Cloning and Regulation of an Endothelial Cell Protein C/Activated Protein C Receptor
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Patrea L. Pabst
 (B) STREET: 2800 One Atlantic Center
 1201 West Peachtree Street
 (C) CITY: Atlanta
 (D) STATE: Georgia
 (E) COUNTRY: USA
 (F) ZIP: 30309-3450

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: Patent In Release #1.0, Version #1.25

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Pabst, Patrea L.
- (B) REGISTRATION NUMBER: 31,284
- (C) REFERENCE/DOCKET NUMBER: OMRF152

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (404) 873-8794
- (B) TELEFAX: (404) 873-8795

(2) INFORMATION FOR SEQ ID NO:1:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1302 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: misc_feature

(B) LOCATION: 1..1302
(D) OTHER INFORMATION: /note= "Nucleotides 25 through 738 encode the Endothelial Cell Protein Receptor of Sequence ID No. 2."
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGGTCCGGA GCCTCAACT CAGGATGTTG ACAACATTGC TGCCGATACT GCTGCTGTCT	60
GGCTGGGCCT TTTGTAGCCA AGAGGCCTCA GATGGCCTCC AAAGACTTCA TATGCTCCAG	120
ATCTCCTACT TCCGCGACCC CTATCACGTG TGGTACCAGG GCAACGCGTC GCTGGGGGGA	180
CACCTAACGC ACGTGTCTGA AGGCCCAGAC ACCAACACCA CGATCATTTCA GCTGCAGCCCC	240
TTGCAGGAGC CCGAGAGCTG GCGGCGCAGG CAGAGTGGCC TGCAGTCCCTA CCTGCTCCAG	300
TTCCACGGCC TCGTGGCCCT GGTGCACGAG GAGCGGACCT TGGCCTTTCC TCTGACCATC	360
CGTGCTTCC TGGGCTGTGA GGTGCTTCCC GAGGGCTCTA GAGCCCATGT CTTCTTCGAA	420
GTGGCTGTGA ATGGGAGCTC CTTTGTGAGT TTCCGGCCGG AGAGAGCCTT GTGGCAGGCA	480
GACACCCAGG TCACCTCCGG AGTGGTCACC TTCACCCCTGC AGCAGCTCAA TGCCTACAAC	540
CGCACTCGGT ATGAACTCGG GGAATTCCCTG GAGGACACCT GTGTGCAGTA TGTGCAGAAA	600
CATATTTCG CGGAAAACAC GAAAGGGAGC CAAACAAGCC GCTCCTACAC TTCGCTGGTC	660
CTGGGCGTCC TGGTGGGCGG TTTCATCATT GCTGGTGTGG CTGTAGGCAT CTTCTCTGTG	720
ACAGGTGGAC GCGCATGTTA ATTACTCTCC AGCCCCGTCA GAAGGGGCTG GATTGATGGA	780
GGCTGGCAAG GGAAAGTTTC AGCTCACTGT GAAGCCAGAC TCCCAACTG AAACACCAGA	840
AGGTTTGGAG TGACAGCTCC TTTCTTCTCC CACATCTGCC CACTGAAGAT TTGAGGGAGG	900
GGAGATGGAG AGGAGAGGTG GACAAAGTAC TTGGTTTGCT AAGAACCCTAA GAACGTGTAT	960

38

GCTTTGCTGA ATTAGTCTGA TAAGTGAATG TTTATCTATC TTTGTGGAAA ACAGATAATG 1020
 GAGTTGGGGC AGGAAGCCTA TGGGCCATCC TCCAAAGACA GACAGAATCA CCTGAGGCGT 1080
 TCAAAAGATA TAACCAATA AACAAATCAT CCACAATCAA AATACAACAT TCAATACTTC 1140
 CAGGTGTGTC AGACTTGGGA TGGGACGCTG ATATAATAGG GTAGAAAGAA GTAACACGAA 1200
 GAAGTGGTGG AATGTAAAA TCCAAGTCAT ATGGCAGTGA TCAATTATTA ATCAATTAAAT 1260
 AATATTAATA AATTCTTAT ATTTAAAAA AAAAAAAAAA AA 1302

(2) INFORMATION FOR SEQ ID NO:2:
 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 238 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (iii) HYPOTHETICAL: NO
- (ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..365

(D) OTHER INFORMATION: /note= "Endothelial Cell Protein Receptor encoded by nucleotides 1 through 1302 of Sequence ID No. 1."

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1..15

(D) OTHER INFORMATION: /note= "Amino acids 1-15 represent a putative signal sequence."

(ix) FEATURE:

- (A) NAME/KEY: Domain
- (B) LOCATION: 211..236

(D) OTHER INFORMATION: /note= "Amino acids 211-236 represent a putative transmembrane domain."

(ix) FEATURE:

- (A) NAME/KEY: Activ -site
- (B) LOCATION: 47..174

(ix) FEATURE:

(B) LOCATION: 17. 186.

(D) OTHER INFORMATION: /NOCC- limiting agent
and 186 reagent extracellular cysteine

residues."

LOCUS	SEQUENCE DESCRIPTION	SEQ ID NO: 2
(xi)		

Met Leu Thr Thr Leu Leu Pro Ile Leu Leu Ser Gly Trp Ala Phe
10 5 10 15

Cys Ser Gln Asp Ala Ser Asp Gly Leu Gln Arg Leu His Met Leu Gln
20 25 30

Ile Ser Tyr Phe Arg Asp Pro Tyr His Val Trp Tyr Gln Gly Asn Ala
25 35 40 45

Ser Leu Gly Gly His Leu Thr His Val Leu Glu Gly Pro Asp Thr Asn
50 55 60

Thr Thr Ile Ile Gln Leu Gln Pro Leu Gln Glu Pro Glu Ser Trp Ala
65 70 75 80

Arg Thr Gln Ser Gly Leu Gln Ser Tyr Leu Leu Gln Phe His Gly Leu
85 90 95

Val Arg Leu Val His Gln Glu Arg Thr Leu Ala Phe Pro Leu Thr Ile
100 105 110

Arg Cys Phe Leu Gly Cys Glu Leu Pro Pro Glu Gly Ser Arg Ala His
115 120 125

Val Phe Phe Phe Glu Val Ala Val Asn Gly Ser Ser Phe Val Ser Phe Arg
130 135 140

Pro Glu Arg Ala Leu Trp Gln Ala Asp Thr Gln Val Thr Ser Gly Val
 145 150 155 160
 Val Thr Phe Thr Leu Gln Gln Leu Asn Ala Tyr Asn Arg Thr Arg Tyr
 165 170 175
 Glu Leu Arg Glu Phe Leu Glu Asp Thr Cys Val Gln Tyr Val Gln Lys
 180 185 190
 His Ile Ser Ala Glu Asn Thr Lys Gly Ser Gln Thr Ser Arg Ser Tyr
 195 200 205
 Thr Ser Leu Val Leu Gly Val Leu Val Gly Gly Phe Ile Ile Ala Gly
 210 215 220
 Val Ala Val Gly Ile Phe Leu Cys Thr Gly Gly Arg Arg Cys
 225 230 235

We claim.

1. An isolated endothelial cell protein C/activated protein C receptor.

2. The receptor of claim 1 encoded by the nucleotide sequence of Sequence ID No. 1 and degenerative sequences thereof and sequence having conservative substitutions, additions or deletions thereof hybridizing to Sequence ID No. 1 under stringent conditions, which encode the receptor.

3. The receptor of claim 1 having the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

4. The receptor of claim 1 expressed on the surface of a non-human cell or a non-endothelial cell.

5. The receptor of claim 1 in soluble form.

6. The receptor of claim 5 lacking at least a portion of the transmembrane region.

7. A nucleotide sequence encoding an endothelial cell protein C/activated protein C receptor.

8. The sequence of claim 7 having the nucleotide sequence of Sequence ID No. 1 or degenerative sequences thereof and sequence having conservative substitutions, additions or deletions thereof and hybridizing under stringent conditions to Sequence ID No. 1.

9. The sequence of claim 7 encoding the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

10. The sequence of claim 7 further comprising an expression vector.

11. The sequence of claim 10 further comprising an expression host.

12. The sequence of claim 10 expressed on the surface of a non-human cell or a non-endothelial cell.
13. The sequence of claim 7 encoding a soluble form of the receptor.
14. The sequence of claim 7 encoding a fragment of the receptor of at least fourteen consecutive nucleotides in length.
15. The sequence of claim 14 labelled with a detectable label.
16. A method for enhancing an inflammatory response involving blocking of protein C or activated protein C binding to an endothelial cell protein C/activated protein C receptor comprising administering to a patient in need of treatment thereof an amount of a compound blocking binding of protein C or activated protein C to the receptor.
17. The method of claim 16 wherein the compound is selected from the group consisting of antibodies and fragments of antibodies to the receptor, nucleic acid sequences inhibiting expression of the receptor, and synthetic or natural compounds other than proteins, peptides or nucleic acid sequences which inhibit binding.
18. A method for inhibiting an inflammatory response involving administration of a compound selected from the group consisting of EPCR or EPCR fragments and substances that upregulate EPCR expression to a patient in need of treatment thereof.
19. An antibody or antibody fragment specifically immunoreactive with a unique epitope of an isolated endothelial cell protein C/activated protein C receptor.
20. The antibody of claim 19 wherein the receptor is encoded by the nucleotide sequence of

Sequence ID No. 1 and degenerative sequences thereof and sequences having conservative substitutions, additions or deletions thereof and hybridizing to Sequence ID No. 1 under stringent conditions.

21. The antibody of claim 19 wherein the receptor has the amino acid sequence of Sequence ID No. 2 or a sequence having conservative substitutions, additions or deletions thereof.

22. A method for screening for a compound which alters the binding of an endothelial receptor protein to protein C or activated protein C comprising providing an assay for binding of protein C or activated protein C to the receptor protein, adding the compound to be tested to the assay, and

determining if the amount of protein C or activated protein C which is bound to the receptor protein is altered as compared to binding in the absence of the compound to be tested.

23. A method for screening patients for abnormal receptor protein activity or function comprising determining the presence of an

endothelial cell surface receptor binding protein C and activated protein C, and comparing the receptor to determine if the quantity present or the function of the receptor is equivalent to that present in normal cells.

24. A method for screening patients for abnormal receptor protein activity or function comprising determining the presence of an endothelial cell surface receptor binding protein C and activated protein C, and comparing the receptor to determine if the quantity present or the function of the receptor is equivalent to that present in normal cells.


1/10
01/92

FIG. 1a




FIG. 1b




FIG. 1c

2/10




FIG. 2a




FIG. 2b

SUBSTITUTE SHEET (RULE 26)




FIG. 2c




FIG. 2d

SUBSTITUTE SHEET (RULE 26)

4/10




FIG. 3a




FIG. 3b

5/10

FIG. 4a

CAGGTCCGGAGCCTCAACTTCAGGATGTTGACAACATTGCTGCCGATACTGCTGCTGTCT 60

M L T T L L P I L L L S 12

GGCTGGGCCTTTGTAGCGAAGACGCCTCAGATGGCCTCCAAGACTTCATATGCTCCAG 120

G W A F C S Q D A S D G L Q R L H M L Q 32

ATCTCCTACTCCGGACCCCTATCAGTGTGGTACCAGGGCAACGGCTCGCTGGGGGGA 180

I S Y F R D P Y H V W Y Q G N A S L G G 52

CACCTAAGCAGGTGCTGGAAGGCCAGACACCAACACGACGATCATTGCTGCAGGCC 240

H L T H V L E G P D T N T T I I Q L Q P 72

TTGCAGGAGCCGAGAGCTGGGGCGGACGAGAGTGGCCTGCAGTCTACCTGCTCCAG 300

L Q E P E S W A R T Q S G L Q S Y L L Q 92

TTCCAGGGCCTGCTGGCCTGGTGACCAAGGAGCGACCTTGGCCTTCCCTGACCATC 360

F H G L V R L V H Q E R T L A F P L T I 112

CGCTGCTTCCTGGGCTGTAGCTGCCCTCCGAGGGCTCTAGAGCCCATGTCTTCTCGAA 420

R C F L G C E L P P E G S R A H V F F E 132

6/10

FIG. 4b

GTGGCTGTGAATGGAGCTCCTTTGTGAGTTCCGGCCGGAGAGAGCCTTGTGGCAGGCA 480
V A V N G S S F V S F R P E R A L W Q A 152

GACACCAGGTACCTCCGGAGTGGTCACCTTCACCCTGCAGCAGCTCAATGCCTACAAC 540
D T Q V T S G V V T F T L Q Q L N A Y N 172

CGCACTCGGTATGAAC TCGGGAATTCCTGGAGGACACCTGTGTGCAGTATGTGCAGAAA 600
R T R Y E L R E F L E D T C V Q Y V Q K 192

CATATTCGGGAAACACGAAAGGGAGGCCAAACAGCCGCTCCTACACTTCGCTGGTC 660
H I S A E N T K G S Q T S R S Y T S L V 212

CTGGCGTCTGGTGGCGGTTTCATTCCTGGTGGTGGCTGTAGGCATCTTCCTGTGC 720
L C V L V G G F I I A G V A V G I F L C 232

ACAGGTGACGGCGATGTTAATTACTCTCCAGCCCGTCAGAGGGGCTGGATTGATGGA 780

T G G R R C * 238

7/10

FIG. 4c

GGCTGGCAAGGAAAGTTTCAGCTCACTGTGAAGCCAGACTCCCCAACTGAACACCAGA 840
 AGGTTTGGAGTGACAGCTCCTTTCTCTCCACATCTGCCCACTGAAGATTGAGGGAGG 900
 GGAGATGGAGAGGAGAGGTGGACAAAGTACTTGGTTTGCTAAGAACCTAAGAACGTGTAT 960
 GCTTTGCTGAATTAGTCTGATAAGTGAATGTTTATCTTCTTTGTGGAAACAGATAATG 1020
 GAGTTGGGGCAGGAAGCCTATGGCCATCTCCAAAGACAGACAGAAATCACCTGAGGCGT 1080
 TCAAAGATATAACCAATAACAAGTCATCCACAATCAAAATACAACATTCAATACTTC 1140
 CAGGTGTGTCAGACTTGGGATGGGACGCTGATATAATAGGGTAGAAAGAAGTAACACGAA 1200
 GAAGTGGTGGAATGTAAATCCAAAGTCATAATGGCAGTGATCAATTATTAAATCAATTAAT 1260
 AATATTAATAAATTTCTTATATTTAAAAAATAAAAAAATAAAAAA 1302




FIG. 5a

[illegible]

EPCR EGPDTN TITIIQLQPLQEPESWARTQSGLSYLLQFHGLVRLVHQER...TLAFPLTI
CCD41 EGPSQNV TIIQLQPLQEPESWERTESGLQIYLTQFESLVKLVYRER...KENVFFPLTV
CD1d SN..DSDTVRS LKPLWSQGTFSDDQWETLQHFIRVYRSSFTRDVKEFAKML..RUSYPLEL
MCD1.2 SN..DSATISFTKPLWSQGLSNQQWEK LQHMFAQVYRVSFTRDIQELVKMMSPKEDYPIEI

```

***          *      *      *      *      *      *      *
****         *      *      *      *      *      *      *
EPCR   RCFLGCEL.P...PEGSRRAHVFFFEAVNGSSFVFSFRPERALWQADTQVTSGVMTFTLQQQL
CCD41   SC$LGCEL.PEEEEEGSEPHVFFDDVAVNGSAFVSFRPKTAJMVWSGSQEPSKAANFTLRQL
CD1d    QVSAGCEVHP....GNASNNFFHVAFQGKDIUSFGTS..WEP.TQEAPLWNLAIQVL
MCD1.2  QLSTGCEMYIP....GNASESFFHVAEQGKYAIVRFGTS..WQR.VLGAPSWLDLPKVL

```

9/10

FIG. 5b

```

**          **      *      *      *      *      *      *
EPCR       NAYNRTRYELREFLEDTCVQVM.....OKHISAENTKGSQTS.....*
CCD41      NAYNRTRYELQEFLLQDTCVEFL.....ENHITTONMKGSQTG.....*
CD1d       NQDKWTRJETVQWLLNGTCTPQFVSGLLESGKSELKRQVKPKAWLSRGPSPGPGRLLLVCHVSG
MCD1.2     NADQGTSATVQTTLLNDTWPFQFARGLLEAGKS DLEKQEKPVAWLSSVPSSAHHGLQLVCHVSG

```

EPCR
CCD41
CD1d
MCD1.2

```

EPCR      ..... RSYTSLVLGVLVGGFIAGVAVGIFLCTGGRRC
CCD41     ..... RSYTSLVLGIIIMCCFIAGVAVGIFMCTSGRGLII
CD1d      QDIVLYWGGSYTSMGLIIAIAVLACLLFLLIVGFTS.RFKRQTSYQGVL
MCD1.2    QDIILYWDARQAPVGLIIVFIVLIMLVVVGAVVYI.WRRRSAYQDIR

```

1141 : 111.11. 1111.11 11111.1.1.1111

1 MLTKFLPLLLLLPGCALCN...SDGSQSLHMLQISYFQDHHHVRHQGNA 47

49 SLGGHLTHVLEGPDTNTTIIQLQPLQEPESWARTQSGLQSYLLQFHGLVR 98

[illegible]

48 SLGKLLTHTLEGPSQNVTLQLQPWQDPESWERTESGLQIYLTQFESLVK 97

99 LVHQR..TLAFPLTIRCFLGCELP...PEGSRAHVFFEVAVNGSSFVSF 143

1 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041

98 LVYRERKENVFFPLTVSCSLGCELPEEEEEGSEPHVFFDVAVNGSAFVSF 147

144RPERALWQADTQVTSGVFTFLQQLNAYNRTRYELREFLEDTCVQYVQKH193

[illegible]

148 RPKTAVWVSGSQEPSKAANFTLKQLNAYNRTRYELQEFLQDTCVEFLENH 197

194 ISAENTKGSQTSRSYTSVLGVLVGGFIIAGVAVGIFLCTGGRRRC 238

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

198 ITTQNMKGSQTGRSYTSLVLGILMGCFIIAGVAVGIFMCTSGRRC 242

FIG. 6

INTERNATIONAL PATENT REPORT

Internal Application No

PCT/US 95/09636

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 A61K39/395 C12N15/11 A61K38/17
C07K16/28 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 42, 21 October 1994 BALTIMORE, MD, USA, pages 26486-26491, K. FUKUDOME ET AL. 'Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor.' see the whole document</p> <p style="text-align: center;">-/--</p>	1-3,7-11

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

2 January 1996

Date of mailing of the international search report

26.01.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL PATENT REPORT

Internal Application No
PCT/US 95/09636

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	CIRCULATION, vol. 90, no. 4 part 2, October 1994 NEW YORK, NY, USA, page I-N K. FUKUDOME ET AL. 'Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor.' see the whole document	1-3,7-11
P,X	CIRCULATION, vol. 90, no. 4 part 2, October 1994 NEW YORK, NY, USA, page I133 K. FUKUDOME ET AL. 'Identification, cloning and regulation of a novel endothelial cell protein C/activated protein C receptor.' see abstract 0707	1-3,7-11
P,X	THROMBOSIS AND HAEMOSTASIS, vol. 72, no. 3, September 1994 STUTTGART, GERMANY, pages 465-474, N. BANGALORE ET AL. 'High affinity binding sites for activated protein C and protein C on cultured human umbilical vein endothelial cells.' see summary see page 472, right column, line 9 - line 50	1-3,7-11
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 10, 10 March 1995 BALTIMORE, MD, USA, pages 5571-5577, K. FUKUDOME ET AL. 'Molecular cloning and expression of murine and bovine endothelial cell protein C/activated protein C receptor (EPCR).' see abstract see figures 1-5	1-4,7-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 95/09636

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-18
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although these claims are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.